

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/29, C07K 14/415, A61K 39/36, G01N 33/68</b>		A1	(11) International Publication Number: <b>WO 96/13589</b>
			(43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/US95/14362		(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).	
(22) International Filing Date: 24 October 1995 (24.10.95)			
(30) Priority Data: 08/330,275 27 October 1994 (27.10.94) US 08/460,039 2 June 1995 (02.06.95) US			
(71) Applicant: IMMULOGIC PHARMACEUTICAL CORPORATION [US/US]; 610 Lincoln Street, Waltham, MA 02154 (US).		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(72) Inventors: KUO, Mei-chang; 5 Cox Road, Winchester, MA 01890 (US). GARMAN, Richard; 21 Fessenden Road, Arlington, MA 02174 (US). GREENSTEIN, Julia; 174 Mount Vernon Street, West Newton, MA 02165 (US). EVANS, Sean; 6 Lincoln Drive, Acton, MA 01720 (US). AMSBERRY, Kent; 51 Cotter Road, Newton, MA 02168 (US). SHAKED, Zeev; 1038 Sierra Street, Berkeley, CA 94707 (US).			
(74) Agents: REMILLARD, Jane, E. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).			
(54) Title: T CELL EPITOPES OF THE MAJOR ALLERGENS FROM AMBROSIA ARTEMISIIFOLIA			
(57) Abstract			
<p>The present invention provides isolated peptides of the major protein allergens of <i>Ambrosia artemisiifolia</i>, or short ragweed pollen. Peptides within the scope of the invention comprise at least one T cell epitope, or preferably at least two T cell epitopes of a protein allergen selected from the allergens Amb a I.1, Amb a I.2, Amb a I.3, Amb a I.4 and Amb a II. Modified peptides having similar or enhanced therapeutic properties as the corresponding, naturally-occurring allergen or portion thereof, but having reduced side effects are disclosed. The invention also provides nucleic acids having sequences encoding peptides of the invention. Methods of treatment or of diagnosis of sensitivity to ragweed pollen allergens in an individual and therapeutic compositions comprising one or more peptides of the invention are also provided including multi-peptide formulations for human therapeutic use.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## T CELL EPITOPES OF THE MAJOR ALLERGENS FROM AMBROSIA ARTEMISIIFOLIA

### Background of the Invention

5        *Ambrosia artemisiifolia* or short ragweed pollen is the major cause of late summer hayfever in North America and Canada and considerable effort has been expended in trying to identify the major allergens produced by this species. *Amb a I* or Antigen E (AgE) has been reported to be the predominant allergen. (King, T.P., et al. *Biochemistry*, 3:458 (1964)). AgE has been characterized and reported to be a nonglycosylated protein of 38kD molecular mass (King, T.P., *Adv. Immunol.* 23:77 (1976); King, T.P., et al., *Arch. Biochem. Biophys.*, 212:127 (1981)). An immunochemically related protein, *Amb a II* (AgK), has been reported to have similar properties (King, T.P., *Adv. Immunol.* 23:77 (1976); King, T.P., *Biochemistry*, 11:367 (1972)). *Amb a I* or AgE can be purified using conventional chromatographic or biochemical techniques. However, it has been reported that due to  
15 cleavage of the 38 kD single-chain precursors by the action of a trypsin-like pollen protease, purification often results in the isolation of two noncovalently associated chains of 26 and 12 kD molecular mass, designated  $\alpha$  and  $\beta$ , respectively (King, T.P., et al., *Arch. Biochem. Biophys.*, 212:127 (1981); King, T.P., et al., *Immunochemistry*, 11:83 (1974)). It has been reported that biochemically purified *Amb a I* has been used as an  
20 immunogen to produce murine monoclonal antibodies (mAb) as well as rabbit polyclonal antisera reactive with both the native (Olson, J.R., and D.G. Klapper, *J. Immunol.*, 136:2109 (1986)) and denatured (Smith, et al., *Molec. Immunol.*, 25:355 (1988)) protein.

Early studies using skin tests with *Amb a I*-depleted pollen extract led to the estimation that at least 90% of the allergenic activity in ragweed pollen can be attributed to  
25 *Amb a I* (King, T.P., et al., *Biochemistry*, 3:458 (1964); King, T.P., et al., *Immunol.*, 23:77 (1976)). Competition experiments using murine mAb to inhibit the binding of *Amb a I* by human IgE in ELISA assays have confirmed that *Amb a I* binds a substantial proportion of human ragweed allergic IgE (Olson, J.R., et al., *J. Immunol.*, 136:2109 (1986)). Recently, three cDNAs encoding proteins with properties of *Amb a I* were cloned and *Amb a I* was  
30 reported to be a family of homologous, but distinguishable sequences (Rafnar, T., et al., *J. Biol. Chem.*, 266:1229 (1991)). Rafnar and co-workers reported that the individual cloned members of the *Amb a I* family, designated *Amb a I.1*, *Amb a I.2*, and *Amb a I.3*, share amino acid sequence homology exceeding 80%. The fourth family member designated *Amb a I.D* (*Amb a I.4*) is disclosed in U.S.S.N. 07/529,951, filed May 29, 1990. The deduced amino  
35 acid sequence of *Amb a II* has been disclosed and reported to share approximately 65% sequence identity with the *Amb a I* multigene family of allergens. (Rogers, B. L., et al., *Journal of Immunology* 147:2547-2552 (1992)).

### **Summary of the Invention**

The present invention provides isolated peptides of the major protein allergens of *Ambrosia artemisiifolia* including peptides derived from the family of related proteins, previously designated *Amb a* IA, *Amb a* IB, *Amb a* IC, and *Amb a* ID. These allergens have  
5 been renamed according to the IUIS approved nomenclature as *Amb a* I.1 (*Amb a* IA), *Amb a* I.2 (*Amb a* IB), *Amb a* I.3 (*Amb a* IC) and *Amb a* I.4 (*Amb a* ID). Peptides within the scope of the invention comprise at least one T cell epitope, preferably at least two T cell epitopes, of a protein allergen selected from the family of *Amb a* I allergens and *Amb a* II. The invention further provides peptides comprising at least two regions, each region comprising  
10 at least one T cell epitope of a ragweed pollen allergen. The regions are derived from the same or from different ragweed pollen allergens.

The invention also provides modified peptides having similar or enhanced therapeutic properties as the corresponding, naturally-occurring allergen or portion thereof, but having reduced side effects as well as modified peptides having improved properties such as  
15 increased solubility and stability. Peptides of the invention are capable of modifying, in a ragweed pollen-sensitive individual to whom they are administered, the allergic response of the individual to a ragweed pollen allergen. Methods of treatment or of diagnosis of sensitivity to a ragweed pollen allergen in an individual and therapeutic compositions comprising one or more peptides of the invention are also provided and human clinical  
20 testing is described.

### **Brief Description of the Drawings**

Fig. 1 shows Western blot analysis of IgE binding to recombinant *Amb a* I proteins.

Fig. 2 is a graphic representation of a direct binding assay of IgE from a single  
25 ragweed allergic patient to recombinant *Amb a* I and *Amb a* II proteins.

Fig. 3 is a graphic representation of the results of a direct binding assay of IgE from pooled human sera to native *Amb a* I, *Amb a* II, recombinant *Amb a* I.1, recombinant *Amb a* II and pollen extract.

Fig. 4A and 4B are graphic representations depicting the responses of lymph node  
30 cells isolated from mice tolerized *in vivo* with either *Amb a* I.1 or PBS and CFA and challenged *in vitro* with various antigens.

Fig. 5A-5F are graphic representations depicting the responses of lymph node cells isolated from mice tolerized *in vivo* with *Amb a* I.1 or pollen extract, challenged with *Amb a* I.1, and tested with various antigens.

Fig. 6A-6F are graphic representations depicting the responses of lymph node cells  
35 isolated from mice tolerized with pollen extract challenged with pollen extract, and tested with various antigens.

Fig. 7 shows various peptides of desired length derived from the *Amb a* I.1, *Amb a* I.2 and *Amb a* I.3 protein allergens.

Fig. 8 is a graphic representation depicting the responses of T cell lines from 39 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to various overlapping *Amb a* I.1 peptides and selected *Amb a* I.2 and *Amb a* I.3 peptides by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 9 shows selected peptides of desired lengths derived from the *Amb a* I.1 protein allergen.

Fig. 10 is a graphic representation depicting the responses of T cell lines from 48 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 1 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 11 is a graphic representation depicting the responses of T cell lines from 48 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 2 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 12 is a graphic representation depicting the responses of T cell lines from 48 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 3 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 13 is a graphic representation depicting the responses of T cell lines from 48 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 4 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 14 shows selected peptides of desired lengths derived from the *Amb a* I.1 protein allergen and the *Amb a* I.3 protein allergen.

Fig. 15 is a graphic representation depicting the responses of T cell lines from 23 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 1 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 16 is a graphic representation depicting the responses of T cell lines from 23 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 2 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 17 is a graphic representation depicting the responses of T cell lines from 23 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 3 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 18 is a graphic representation depicting the responses of T cell lines from 23 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 4 of the *Amb a* I.1 protein, by percent of positive responses within the individuals tested, the mean stimulation index of positive responses for the peptide and the ranked sum of peptide responses.

Fig. 19 is a graphic representation depicting the responses of T cell lines of 9 patients primed *in vitro* to recombinant *Amb a* I.1 protein or recombinant *Amb a* I.3 protein and analyzed for response to selected peptides derived from *Amb a* I.1, by percent of positive responses within the individuals tested and the mean stimulation index of positive responses for the peptide.

Fig. 20 is a graphic representation depicting the responses of T cell lines of 9 patients primed *in vitro* to recombinant *Amb a* I.1 protein or recombinant *Amb a* I.3 protein and analyzed for response to selected peptides derived from *Amb a* I.3, by percent of positive responses within the individuals tested and the mean stimulation index of positive responses for the peptide.

Fig. 21 is a graphic representation of a direct binding assay of IgE from a single ragweed allergic patient to peptides derived from *Amb a* I.

Fig. 22 is a graphic representation depicting the responses of T cell lines of 28 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from *Amb a* I.1 by percent of positive responses within the individuals tested, the mean stimulation index of positive responses to the peptide and the ranked sum of peptide responses.

Fig. 23 is a graphic representation depicting the responses of T cell lines of 28 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to selected peptides derived from Region 4 of *Amb a* I.1 by percent of positive responses within the individuals tested, the mean stimulation index of positive responses to the peptide and the ranked sum of peptide responses.

Fig. 24 is a graphic representation depicting the responses of T cell lines of 32 patients primed *in vitro* to recombinant *Amb a* I.1 protein and analyzed for response to six selected peptides derived from *Amb a* I.1 by percent of positive responses within the individuals tested, the mean stimulation index of positive responses to the peptide and the ranked sum of peptide responses.

Fig. 25 shows various peptides derived from peptide RAE 70.1 which include modifications designed to increase the solubility of the peptide.

Fig. 26 is a graphic representation depicting the response of a T cell line from patient 956.2 primed in vitro to *Fel d 1* and analyzed for response to various peptides derived from the *Amb a I.1* protein.

Fig. 27 is a graphic representation depicting the response of a T cell line from patient 119 primed in vitro with recombinant *Amb a I.1* and analyzed for response to various modified peptides derived from Region 2 of the *Amb a I.1* protein by tritiated thymidine incorporation.

Fig. 28 is a graphic representation depicting the response of a T cell line from patient 1199 primed in vitro with recombinant *Amb a I.1* and analyzed for response to various modified peptides derived from Region 2 of the *Amb a I.1* protein by tritiated thymidine incorporation.

Fig. 29 is a graphic representation depicting the response of a T cell clone generated by limiting dilution from an *Amb a I.1* specific T cell line stimulated with the AMB 2-10.1 peptide, primed in vitro w/recombinant *Amb a I.1* and analyzed for response to various modified peptides derived from Region 2 of the *Amb a I.1* protein by tritiated thymidine incorporation.

Fig. 30 is a graphic representation depicting the percent of total histamine release in blood samples from 8 ragweed-allergic patients in response to selected peptides derived from the *Amb a I.1* protein.

Fig. 31 is a graphic representation depicting the responses of T cell lines from 39 patients primed in vitro to purified native *Amb a I.1* protein and analyzed for response to various overlapping *Amb a I.1* peptides by percent of positive responses within the individuals tested (above the bar), the mean stimulation index of positive responses for the peptide (in parenthesis above the bar) and the ranked sum of peptide responses (X axis).

### Detailed Description of the Invention

The present invention provides isolated peptides derived from the major protein allergens of *Ambrosia artemisiifolia*. As used herein, a peptide refers to an amino acid sequence having fewer amino acids than the entire amino acid sequence of a protein from which the peptide is derived. Peptides of the invention include peptides derived from *Amb a I.1*, *Amb a I.2*, *Amb a I.3*, *Amb a I.4* and *Amb a II* which comprise at least one T cell epitope of the allergen.

Peptides comprising at least two regions, each region comprising at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia* are also within the scope of the invention. Each region of such peptides is derived from the same or from different ragweed pollen allergens. Isolated peptides or regions of isolated peptides, each comprising at least two T cell epitopes of a ragweed pollen allergen are particularly desirable for increased therapeutic effectiveness. Peptides which are immunologically related (e.g., by antibody or T cell cross-reactivity) to peptides of the present invention are also within the scope of the

invention. Peptides immunologically related by antibody cross-reactivity are bound by antibodies specific for a peptide of a protein allergen of *Ambrosia artemisiifolia*. Peptides immunologically related by T cell cross-reactivity are capable of reacting with the same T cells as a peptide of the invention.

5       The present invention also pertains to a ragweed pollen allergen encoded by a nucleic acid sequence of clone IPC1/5. The full-length nucleic acid sequence of clone IPC1/5 has been determined and the encoded protein has been produced recombinantly in both the pSEM vector (as a fusion protein with  $\beta$ -galactosidase) and the pET11d vector. The recombinant protein was determined to bind approximately 10-20% of allergic serum IgE on a Western blot. The protein encoded by clone IPC1/5 was found to have a high degree of amino acid sequence homology with cysteine proteinase inhibitors in man and rice. The protein has 10       66.6% homology with the rice protein oryzacystatin-I. The nucleic acid sequence and deduced amino acid sequence of the allergen encoded by clone IPC1/5 is represented in SEQ ID NO. 11 and 12.

15       Isolated proteins and isolated peptides of the invention can be produced by recombinant DNA techniques in a host cell transformed with a nucleic acid having a sequence encoding such protein or peptide. The isolated proteins and isolated peptides of the invention can also be produced by chemical synthesis. In certain limited situations, isolated peptides can be produced by chemical cleavage of a protein allergen. When a protein or 20       peptide is produced by recombinant techniques, host cells transformed with a nucleic acid having a sequence encoding the protein or peptide or the functional equivalent of the nucleic acid sequence are cultured in a medium suitable for the cells and protein or peptides can be purified from cell culture medium, host cells, or both using techniques known in the art for purifying proteins and peptides including ion-exchange chromatography, gel filtration 25       chromatography, ultrafiltration, electrophoresis or immunopurification with antibodies specific for the protein or peptide, the protein allergen of *Ambrosia artemisiifolia* from which the peptide is derived, or a portion thereof. By isolated is meant that protein and peptides of the present invention are substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or substantially free of chemical precursors or 30       other chemicals when synthesized chemically. Recombinant ragweed pollen proteins including recombinant *Amb a* I.1, *Amb a* I.2, *Amb a* I.3, *Amb a* I.4, and *Amb a* II have been produced.

      Suitable expression vectors for producing recombinant protein and recombinant peptides of the invention include pTRC, pGEX, pMAL, pRIT5, pET11d and pCA. The use 35       of pTRC, pET11d and pGEX as expression vectors will result in expression of ragweed pollen protein as an unfused protein. The use of pMAL, pRIT5, pCA and pSEM as expression vectors will result in expression of ragweed pollen protein fused to maltose E binding protein (pMAL), protein A (pRIT5), truncated protein A (pCA), or  $\beta$ -galactosidase (pSEM). Suitable expression vectors are commercially available. When produced as a



fusion protein, recombinant ragweed pollen protein can be recovered from the fusion protein through enzymatic or chemical (e.g., cyanogen bromide or dilute acid) cleavage and biochemical purification. For example, enzymatic cleavage sites for Factor Xa or thrombin can be introduced at the fusion junction between the carrier protein (e.g., Protein A) and the ragweed pollen protein. Suitable host cells for expression of recombinant ragweed pollen protein include bacteria, yeast and insect or mammalian cells. Appropriate vectors for expression in yeast include YepSec, pMFa and JRY88. These vectors are also commercially available.

To obtain isolated peptides of the present invention, a ragweed pollen allergen is divided into non-overlapping peptides of desired lengths or overlapping peptides of desired lengths as discussed in Example V which may be produced recombinantly, synthetically or in certain limited situations by chemical cleavage of the allergen. Peptides comprising at least one T cell epitope are capable of reducing T cell responsiveness or inducing T cell nonresponsiveness. To determine peptides comprising at least one T cell epitope, isolated peptides are tested by, for example, T cell biology techniques to determine whether the peptides elicit a T cell response or induce T cell non responsiveness. Those peptides found to elicit a T cell response or induce T cell nonresponsiveness are defined as having T cell stimulating activity. As discussed in the Examples, human T cell stimulating activity can be tested by culturing T cells obtained from an individual sensitive to a ragweed pollen allergen, (i.e., an individual who has an IgE mediated immune response to a ragweed pollen allergen) with a peptide derived from the allergen and determining whether proliferation of T cells occurs in response to the peptide as measured, e.g., by cellular uptake of tritiated thymidine. As described in detail in the Examples, stimulation indices for responses by T cells to peptides can be calculated as the maximum CPM in response to a peptide divided by the medium control CPM. As used throughout this application, a peptide comprising at least one T cell epitope, when determined by T cell stimulation requires a stimulation index of at least 2.0. A peptide having a T cell stimulation index of 2.0 is considered useful as a therapeutic agent. Preferred peptides have a stimulation index of at least 2.5, more preferably at least 3.5, and most preferably at least 5.0.

In order to determine precise T cell epitopes by, for example, fine mapping techniques, a peptide having T cell stimulating activity and thus comprising at least one T cell epitope as determined by T cell biology techniques is modified by addition or deletion of amino acid residues at either the amino or carboxy terminus of the peptide and tested to determine a change in T cell reactivity to the modified peptide. If two or more peptides which share an area of overlap in the native protein sequence are found to have human T cell stimulating activity, as determined by T cell biology techniques, additional peptides can be produced comprising all or a portion of such peptides and these additional peptides can be tested by a similar procedure. Following this technique, peptides are selected and produced recombinantly or synthetically. Peptides are selected based on various factors, including the

strength of the T cell response to the peptide (e.g., stimulation index), the frequency of the T cell response to the peptide in a population of individuals sensitive to ragweed pollen, and the potential cross-reactivity of the peptide with *Amb a I* family members and *Amb a II*. The physical and chemical properties of these selected peptides (e.g., solubility, stability) are examined to determine whether the peptides are suitable for use in therapeutic compositions or whether the peptides require modification as described herein. The ability of the selected peptides or selected modified peptides to stimulate human T cells (e.g., induce proliferation, lymphokine secretion) is determined.

— In addition, preferred peptides of the invention do not bind immunoglobulin E (IgE) or bind IgE to a substantially lesser extent (i.e. preferably at least 100 fold and more preferably at least 1000 fold less) than the protein allergen from which the peptide is derived binds IgE. Recombinant ragweed pollen allergens including recombinant *Amb a I.1*, *Amb a I.2*, *Amb a I.3*, *Amb a I.4*, and *Amb a II* have been produced and shown to have reduced IgE binding activity as compared to the corresponding native protein allergen (See Fig. 3). The major complications of standard immunotherapy are IgE-mediated responses such as anaphylaxis. Immunoglobulin E is a mediator of anaphylactic reactions which result from the binding and cross-linking of antigen to IgE on mast cells or basophils and the release of mediators (e.g., histamine, serotonin, eosinophil chemotactic factors). Thus, anaphylaxis in a substantial percentage of a population of individuals sensitive to ragweed pollen allergen could be avoided by the use in immunotherapy of a recombinant protein, a peptide or peptides which do not bind IgE in a substantial percentage (e.g., at least about 75%) of a population of individuals sensitive to ragweed pollen allergen, or if the protein or peptide(s) binds IgE, such binding does not result in the release of mediators from mast cells or basophils. Additionally, the risk of anaphylaxis can be reduced by the use in immunotherapy of a recombinant protein, a peptide or peptides which have reduced IgE binding. Moreover, recombinant protein or peptides which have minimal IgE stimulating activity are desirable for therapeutic effectiveness. Minimal IgE stimulating activity refers to IgE production that is less than the amount of IgE production and/or IL-4 production stimulated by the native protein allergen (e.g., *Amb a I.1*).

A peptide or recombinant protein of the invention, when administered to a ragweed pollen-sensitive individual, is capable of modifying the allergic response of the individual to the allergen. Particularly, peptides of the invention comprising at least one T cell epitope of a ragweed pollen allergen or at least two regions derived from a ragweed pollen allergen each comprising at least one T cell epitope, when administered to a ragweed pollen-sensitive individual are capable of modifying the T cell response of the individual to the allergen. As used herein, modification of the allergic response of a ragweed pollen-sensitive individual to a ragweed pollen allergen can be defined as non-responsiveness or diminution in symptoms to a ragweed pollen allergen, as determined by standard clinical procedures (see e.g., Varney et al., British Medical Journal 302: 265-269 (1990)).

As a result of the work described herein, peptides derived from ragweed pollen allergens comprising at least one T cell epitope have been produced. T cell epitopes are believed to be involved in initiation and perpetuation of the immune response to ragweed pollen allergen(s) which are responsible for the clinical symptoms of ragweed pollen allergy.

5 These T cell epitopes are thought to trigger early events at the level of the T helper cell by binding to an appropriate HLA molecule on the surface of an antigen presenting cell and stimulating the relevant T cell subpopulation. These events lead to T cell proliferation, lymphokine secretion, local inflammatory reactions, the recruitment of additional immune

10 cells to the site, and activation of the B cell cascade leading to production of antibodies. One isotype of these antibodies, IgE, is fundamentally important in the development of allergic symptoms and its production is influenced early in the cascade of events, at the level of the T helper cell, by the nature of the lymphokines secreted. A T cell epitope is the basic element or smallest unit of recognition by a T cell receptor, where the epitope comprises amino acid

15 residues essential to receptor recognition which may be contiguous and/or non-contiguous in the amino acid sequence of the protein. Amino acid sequences which mimic those of T cell epitopes and which modify the allergic response to protein allergens of *Ambrosia artemisiifolia* are within the scope of this invention.

Exposure of ragweed pollen allergic patients to peptides of the present invention, in a non-immunogenic form, may induce T cell non-responsiveness of appropriate T cell

20 subpopulations such that they become non-responsive to ragweed pollen allergen(s) and do not participate in mounting an immune response upon such exposure.

While not intending to be limited to any theory, it is believed that T cell non-responsiveness (which includes reduced T cell responsiveness) is induced as a result of not providing an appropriate costimulatory signal sometimes referred to as a "second signal"

25 Briefly, it is believed that stimulation of T cells requires two types of signals. the first is the recognition by the T cell via the T cell receptor of appropriate MHC-associated processed antigens on antigen presenting cells (APCs) and the second type of signal is referred to as a costimulatory signal(s) or "second signal" which may be provided by certain competent APCs. When a composition of the invention is administered without

30 adjuvant, it is believed that competent APCs which are capable of producing the second signal or costimulatory signal are not engaged in the stimulation of appropriate T cells therefore resulting in T cell nonresponsiveness or reduced T cell responsiveness. In addition, there are a number of antibodies or other reagents capable of blocking the delivery of costimulatory signals such as the "second signal" which include, but are not

35 limited to B7 (including B7-1, B7-2, and BB-1), CD28, CTLA4, CD40 CD40L CD54 and CD11a/18 (Jenkins and Johnson, *Current Opinion in Immunology*, 5:361-367 (1993), and Clark and Ledbetter, *Nature*, 367:425-428 (1994)) Thus, a peptide of the invention may be administered in nonimmunogenic form as discussed above, in conjunction with a reagent

capable of blocking costimulatory signals such that the level of T cell nonresponsiveness is enhanced.

In addition, administration of a peptide of the present invention may modify the lymphokine secretion profile as compared with exposure to the naturally-occurring ragweed pollen allergen or portion thereof (e.g., result in a decrease of IL-4 and/or an increase in IL-2). Furthermore, exposure to a peptide of the invention may influence T cell subpopulations which normally participate in the response to ragweed pollen allergen(s) such that these T cells are drawn away from the site(s) of normal exposure to the allergen (e.g., nasal mucosa, skin, and lung) towards the site(s) of therapeutic administration of the peptide. This redistribution of T cell subpopulations may ameliorate or reduce the ability of an individual's immune system to stimulate the immune response at the site of normal exposure to the ragweed pollen allergen(s), resulting in a diminution in allergic symptoms.

Isolated peptides of the invention comprise at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia* (i.e., the peptide comprises at least approximately seven amino acid residues of the protein allergen). For purposes of therapeutic effectiveness, therapeutic compositions of the invention preferably comprise at least two T cell epitopes of a ragweed pollen allergen. Accordingly, isolated peptides of the invention preferably comprise at least two T cell epitopes (i.e., the peptide comprises at least approximately eight amino acid residues, and preferably fifteen amino acid residues). Additionally, isolated peptides of the invention preferably comprise a sufficient percentage of the T cell epitopes of the entire protein allergen such that upon administration of the peptide to an individual sensitive to ragweed pollen, T cells of the individual are rendered non-responsive to the protein allergen. Isolated peptides of the invention comprising up to approximately 45 amino acid residues in length, and most preferably up to approximately 30 amino acid residues in length are particularly desirable as increases in length may result in difficulty in peptide synthesis as well as retention of an undesirable property (e.g., immunoglobulin binding or enzymatic activity) due to maintenance of conformational similarity between the peptide and the protein allergen from which it is derived. All of the peptides shown in Fig. 8 were found to have human T cell stimulating activity.

To determine whether a peptide (candidate peptide) or a combination of candidate peptides are likely contain a sufficient percentage of T cell epitopes of ragweed protein antigen to induce non-responsiveness in a substantial percentage of a population of individuals sensitive to the protein antigen, an algorithm can be used. In accordance with one such algorithm, a human T cell stimulation index (discussed above) for the peptide(s) in an *in vitro* T cell proliferation assay is calculated for each individual tested in a population of individuals sensitive to ragweed protein allergen. The remaining peptides in the *in vitro* T cell proliferation assay are overlapping peptides (overlapping by between about 5 - 10 amino acid residues) which cover the remainder of the protein not covered by the candidate peptide(s), which peptides are at least 12 amino acids long and which are preferably no longer than 30

and more preferably no longer than 25 amino acid residues in length. A human T cell stimulation index for each such remaining peptide in the set of peptides produced in the *in vitro* T-cell proliferation assay with T-cells obtained from each individual in the population of individuals tested is calculated and added together. For each individual, the human T cell stimulation index for the candidate peptide(s) is divided by the sum of the human T cell stimulation indices of the remaining peptides in the set of peptides tested to determine a percent. This percent is obtained for at least twenty (20) and preferably at least thirty (30) individuals sensitive to the protein antigen of interest and a mean percent is determined (the percentage of positive T cell responses (S.I. greater than or equal to 2.0) in response to the candidate peptide or combination of candidate peptides). A mean percent of about 10% or greater for the candidate peptide(s) together with a percent positive of at least about 60%, preferably about 75% and more preferably about 90%, or most preferably 100%, indicates that the candidate peptide(s) selected is likely to contain a sufficient percentage of T cell epitopes to induce T cell non responsiveness in a substantial percentage of a population of individuals sensitive to TRFP.

Preferred peptides comprise all or a portion of the areas of major T cell reactivity within the *Amb a I.1* protein allergen, i.e., Region 1, Region 2, Region 3 and Region 4. Each area is broadly defined as follows: Region 1 comprises amino acid residues 48-107; Region 2 comprises amino acid residues 171-216; Region 3 comprises amino acid residues 278-322; and Region 4 comprises amino acid residues 331-377. Preferred areas of major T cell reactivity within each Region comprise: amino acid residues 57-101; amino acid residues 182-216; amino acid residues 280-322; and amino acid residues 342-377. Similar areas of major T cell reactivity can be found within the other *Amb a I* family members (i.e., *Amb a I.2*, *Amb a I.3* and *Amb a I.4*), and *Amb a II*. As shown in Example VIII, the *Amb a I* protein allergens and *Amb a II* demonstrate a high degree of T cell cross-reactivity. Given this cross-reactivity, shared areas of major T cell reactivity and shared T cell epitopes are likely to be found in conserved regions between *Amb a I* and the remaining *Amb a I* family members and *Amb a II*. For example, *Amb a I.1* stimulated T cells have been shown to recognize both *Amb a I.1* derived peptides and homologous *Amb a I.3* derived peptides (See Example IX). Similarly, *Amb a I.3* stimulated T cell recognize both *Amb a I.1* and *Amb a I.3* derived peptides.

Preferred ragweed pollen peptides comprise all or a portion of the following peptides: RAE 67.1 (SEQ ID NO:13); RAE 57.1 (SEQ ID NO:14); RAE 24.E (SEQ ID NO:15); RAE 24.1 (SEQ ID NO:16); RAE 22.E (SEQ ID NO:17); RAE 22.E-1 (SEQ ID NO:18); RAE 3.D (SEQ ID NO:19); RAE 3.1 (SEQ ID NO:20); RAE 22.E-2 (SEQ ID NO:21); RAE 5.D (SEQ ID NO:22); RAE 6.D (SEQ ID NO:23); RAE 6.1 (SEQ ID NO:24); RAE 7.D (SEQ ID NO:25); RAE 7.D-1 (SEQ ID NO:26); RAE 40.1-6 (SEQ ID NO:27); RAE 40.1-5 (SEQ ID NO:28); RAE 40.1-4 (SEQ ID NO:29); RAE 40.D (SEQ ID NO:30); RAE 40.1 (SEQ ID NO:31); RAE 61.1 (SEQ ID NO:32); RAE 80.1 (SEQ ID NO:33); RAE 45.1 (SEQ ID

NO:34); RAE 75.1 (SEQ ID NO:35); RAE 62.1 (SEQ ID NO:36); RAE 69.1 (SEQ ID NO:37); RAE 69.1-1 (SEQ ID NO:38); RAE 69.1-2 (SEQ ID NO:39); RAE 69.1-3 (SEQ ID NO:40); RAE 70.1-3 (SEQ ID NO:41); RAE 70.1-2 (SEQ ID NO:42); RAE 70.1-1 (SEQ ID NO:43); RAE 70.1 (SEQ ID NO:44); RAE 71.1 (SEQ ID NO:45); RAE 65.1 (SEQ ID NO:46); RAE 63.1 (SEQ ID NO:47); RAE 76.1 (SEQ ID NO:48); RAE 27.1 (SEQ ID NO:49); RAE 66.1 (SEQ ID NO:50); RAE 66.1-1 (SEQ ID NO:51); RAE 66.1-2 (SEQ ID NO:52); RAE 66.1-3 (SEQ ID NO:53); RAE 64.1-3 (SEQ ID NO:54); RAE 64.1-2 (SEQ ID NO:55); RAE 64.1-1 (SEQ ID NO:56); RAE 64.1 (SEQ ID NO:57); RAE 73.1 (SEQ ID NO:58); RAE 74.1 (SEQ ID NO:59); RAE 74.1-1 (SEQ ID NO:60); RAE 29.1 (SEQ ID NO:61); RAE 29.1-1 (SEQ ID NO:62); RAE 28+29 (SEQ ID NO:63); RAE 29.1-2 (SEQ ID NO:64); RAE 29.1-3 (SEQ ID NO:65); RAE 29.1-4 (SEQ ID NO:66); RAE 28.1-3 (SEQ ID NO:67); RAE 28.1-2 (SEQ ID NO:68); RAE 28.1-1 (SEQ ID NO:69); RAE 28.1 (SEQ ID NO:70); RAE 20.1 (SEQ ID NO:71); RAE 20.1-3 (SEQ ID NO:72); RAE 20.1-2 (SEQ ID NO:73); RAE 20.1-1 (SEQ ID NO:74); RAE 21.1 (SEQ ID NO:75); RAE 17.1 (SEQ ID NO:76); RAE 55.1 (SEQ ID NO:77); RAE 76.6 (SEQ ID NO:78); RAE 67.15 (SEQ ID NO:79); RAE 45.15 (SEQ ID NO:80); RAE 27.15 (SEQ ID NO:81); AMB 1-1.1 (SEQ ID NO:85); AMB 1-2.1 (SEQ ID NO:86); AMB 1-3.1 (SEQ ID NO:87); AMB 1-4.1 (SEQ ID NO:84); AMB 1-5.1 (SEQ ID NO:83); AMB 1-6.1 (SEQ ID NO:82); AMB 1-4.15 (SEQ ID NO:88); AMB 1-2.15 (SEQ ID NO:89); AMB 2-4.1 (SEQ ID NO:90); AMB 2-3.1 (SEQ ID NO:91); AMB 2-5.1 (SEQ ID NO:92); AMB 2-6.1 (SEQ ID NO:93); AMB 2-2.1 (SEQ ID NO:94); AMB 2-1.1 (SEQ ID NO:95); AMB 2-7.1 (SEQ ID NO:96); AMB 2-8.1 (SEQ ID NO:97); AMB 2-9.1 (SEQ ID NO:98); AMB 2-10.1 (SEQ ID NO:99); AMB 2-11.1 (SEQ ID NO:100); AMB 2-1.15 (SEQ ID NO:101); AMB 3-4.1 (SEQ ID NO:103); AMB 3-5.1 (SEQ ID NO:102); AMB 3-3.1 (SEQ ID NO:104); AMB 3-2.1 (SEQ ID NO:105); AMB 3-1.1 (SEQ ID NO:106); AMB 3-4.15 (SEQ ID NO:107); AMB 3-1.15 (SEQ ID NO:108); AMB 4-8.1 (SEQ ID NO:109); AMB 4-9.1 (SEQ ID NO:110); AMB 4-6.1 (SEQ ID NO:111); AMB 4-5.1 (SEQ ID NO:112); AMB 4-3.1 (SEQ ID NO:113); AMB 4-2.1 (SEQ ID NO:114); AMB 4-1.1 (SEQ ID NO:115); AMB 4-3.15 (SEQ ID NO:116); Amb 2-18.1 (SEQ ID NO:126); Amb 2-19.1 (SEQ ID NO:127); Amb 2-20.1 (SEQ ID NO:128); Amb 2-21.1 (SEQ ID NO:129); Amb 2-22.1 (SEQ ID NO:130); Amb 2-23.1 (SEQ ID NO:131); Amb 2-26.1 (SEQ ID NO:132); Amb 28.1 (SEQ ID NO:133); Amb 2-30.1 (SEQ ID NO:134); Amb 2-32.1 (SEQ ID NO:135); Amb 2-33.1 (SEQ ID NO:136); Amb 2-34.1 (SEQ ID NO:137); Amb 2-35.1 (SEQ ID NO:138); Amb 2-36.1 (SEQ ID NO:139); Amb 2-37.1 (SEQ ID NO:140); Amb 2-38.1 (SEQ ID NO:141); AMB 4-9.1EP (SEQ ID NO:142); AMB 4-9.1NP (SEQ ID NO:143); AMB 4-9.1AP (SEQ ID NO:144); AMB 4-9.1SP (SEQ ID NO:145); AMB 4-9.1QP (SEQ ID NO:146); AMB 4-9.1DA (SEQ ID NO:147); amb 4-9.1DS (SEQ ID NO:148); AMB 4-9.1DG (SEQ ID NO:149); and RA-02.1 (SEQ ID NO:150), the amino acid sequences of such peptides being shown in Figs. 7, 14, 23, 24 and 25. Particularly preferred peptides comprise all or a portion of the following peptides: AMB 1-2.1 (SEQ ID NO:86);

AMB 2-6.1 (SEQ ID NO:93); AMB 2-4.1 (SEQ ID NO:90); Amb 2-36.1 (SEQ ID NO:139); Amb 2-38.1 (SEQ ID NO:141); RA-02.1 (SEQ ID NO:150); AMB 2-9.1 (SEQ ID NO:98); AMB 3-5.1 (SEQ ID NO:102); and AMB 4-9.1 (SEQ ID NO:110).

Another embodiment of the present invention provides peptides comprising at least  
5 two regions, each region comprising at least one T cell epitope of a protein allergen of  
*Ambrosia artemisiifolia* (e.g., each region comprises at least approximately seven amino acid  
residues). These peptides comprising at least two regions can comprise as many amino acid  
residues as desired and preferably comprise at least about 7, more preferably at least about  
15, even more preferably about 30 and most preferably at least about 40 amino acid residues  
10 of a ragweed pollen allergen. Each region of such peptide preferably comprises up to 45  
amino acid residues in length, more preferably up to 40 residues in length and most  
preferably up to 30 amino acid residues in length as increases in length of a region may result  
in difficulty in peptide synthesis as well as retention of an undesirable property (e.g.,  
immunoglobulin binding or enzymatic activity) due to maintenance of conformational  
15 similarity between the peptide and the protein allergen from which it is derived. If desired,  
the amino acid sequences of the regions can be produced and joined by a linker to increase  
sensitivity to processing by antigen-presenting cells. Such linker can be any non-epitope  
amino acid sequence or other appropriate linking or joining agent. To obtain preferred  
peptides comprising at least two regions, each comprising at least one T cell epitope, the  
20 regions are arranged in a configuration different from a naturally-occurring configuration of  
the regions in the allergen. For example, the regions containing T cell epitope(s) can be  
arranged in a noncontiguous configuration and can preferably be derived from the same  
protein allergen. Noncontiguous is defined as an arrangement of regions containing T cell  
epitope(s) which is different than that of an amino acid sequence present in the protein  
25 allergen from which the regions are derived. Furthermore, the noncontiguous regions  
containing T cell epitopes can be arranged in a nonsequential order (e.g., in an order different  
from the order of the amino acids of the native protein allergen from which the region  
containing T cell epitope(s) are derived in which amino acids are arranged from an amino  
terminus to a carboxy terminus). A peptide can comprise at least 15%, at least 30%, at least  
30 50% or up to 100% of the T cell epitopes of a ragweed pollen allergen.

The individual peptide regions can be produced and tested to determine which regions  
bind immunoglobulin E specific for a ragweed pollen allergen and which of such regions  
would cause the release of mediators (e.g., histamine) from mast cells or basophils. Those  
peptide regions found to bind immunoglobulin E and cause the release of mediators from  
35 mast cells or basophils in greater than approximately 10-15% of the allergic sera tested are  
preferably not included in the peptide regions arranged to form peptides of the invention.

Preferred peptides of the invention comprise two or more regions derived from the  
same or from different ragweed pollen allergens (e.g., *Amb a I.1*, *Amb a I.2*, *Amb a I.3*, *Amb a I.4* and *Amb a II*). For example, one region can be derived from *Amb a I.1* and one region

can be derived from *Amb a* I.2; one region can be derived from *Amb a* I.1 and one region can be derived from *Amb a* I.3; one region can be derived from *Amb a* I.1 and one region can be derived from *Amb a* I.4; one region can be derived from *Amb a* I.2 and one region can be derived from *Amb a* I.3; one region can be derived from *Amb a* I.2 and one region can be derived from *Amb a* I.4; one region can be derived from *Amb a* I.3 and one region can be derived from *Amb a* I.4; one region can be derived from *Amb a* I.1 and one region can be derived from *Amb a* II; one region can be derived from *Amb a* I.2 and one region can be derived from *Amb a* II; one region can be derived from *Amb a* I.3 and one region can be derived from *Amb a* II; and one region can be derived from *Amb a* I.4 and one region can be derived from *Amb a* II. Alternatively, the regions can be derived from the same protein allergen, e.g., *Amb a* I.1 and *Amb a* I.1, etc.

Regions of a peptide of the invention preferably comprise all or a portion of Region 1, Region 2, Region 3 and Region 4 of *Amb a* I.1, and the above discussed preferred areas of major T cell reactivity within each Region. If Region 1, 2, 3 or 4 is found to bind IgE and cause the release of mediators from mast cells or basophils, then it is preferred that more than one region of the peptide comprise such Region and that the various regions of the peptide do not bind IgE or cause release of mediators from mast cells or basophils. Examples of preferred regions include: AMB 1-1.1 (SEQ ID NO:85); AMB 1-2.1 (SEQ ID NO:86); AMB 1-3.1 (SEQ ID NO:87); AMB 1-4.1 (SEQ ID NO:84); AMB 1-5.1 (SEQ ID NO:83); AMB 1-6.1 (SEQ ID NO:82); AMB 1-4.15 (SEQ ID NO:88); AMB 1-2.15 (SEQ ID NO:89); AMB 2-4.1 (SEQ ID NO:90); AMB 2-3.1 (SEQ ID NO:91); AMB 2-5.1 (SEQ ID NO:92); AMB 2-6.1 (SEQ ID NO:93); AMB 2-2.1 (SEQ ID NO:94); AMB 2-1.1 (SEQ ID NO:95); AMB 2-7.1 (SEQ ID NO:96); AMB 2-8.1 (SEQ ID NO:97); AMB 2-9.1 (SEQ ID NO:98); AMB 2-10.1 (SEQ ID NO:99); AMB 2-11.1 (SEQ ID NO:100); AMB 2-1.15 (SEQ ID NO:101); AMB 3-4.1 (SEQ ID NO:103); AMB 3-5.1 (SEQ ID NO:102); AMB 3-3.1 (SEQ ID NO:104); AMB 3-2.1 (SEQ ID NO:105); AMB 3-1.1 (SEQ ID NO:106); AMB 3-4.15 (SEQ ID NO:107); AMB 3-1.15 (SEQ ID NO:108); AMB 4-8.1 (SEQ ID NO:109); AMB 4-9.1 (SEQ ID NO:110); AMB 4-6.1 (SEQ ID NO:111); AMB 4-5.1 (SEQ ID NO:112); AMB 4-3.1 (SEQ ID NO:113); AMB 4-2.1 (SEQ ID NO:114); AMB 4-1.1 (SEQ ID NO:115); AMB 4-3.15 (SEQ ID NO:116); RA-02.1 (SEQ ID NO:150); Amb 2-36.1 (SEQ ID NO:139); and Amb 2-38.1 (SEQ ID NO:141), the amino acid sequences of such regions being shown in Fig. 14, Fig. 24 or Fig. 25, or portions of said regions comprising at least one T cell epitope.

Preferred peptides comprise various combinations of two or more regions, each region comprising all or a portion of Region 1, Region 2, Region 3 or Region 4 of *Amb a* I.1. Preferred peptides comprise various combinations of two or more regions, each region having an amino acid sequence as shown in Fig. 14, such combination of regions including the following: AMB 4-6.1 and RAE 70.1 (SEQ ID NO:111 and SEQ ID NO:44); AMB 4-6.1 and AMB 2-5.1 (SEQ ID NO:111 and SEQ ID NO:92); AMB 4-9.1 and AMB 2-5.1 (SEQ ID NO:110 and SEQ ID NO:92); AMB 4-9.1 and RAE 70.1 (SEQ ID NO:110 and SEQ ID NO:44).



NO:44); AMB 4-6.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:92 and SEQ ID NO:86); AMB 4-9.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:92 and SEQ ID NO:86); AMB 4-6.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:44 and SEQ ID NO:86); AMB 4-9.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:44 and SEQ ID NO:86); AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103); AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:101, and SEQ ID NO:116); AMB 1-2.15, AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:89, SEQ ID NO:101, and SEQ ID NO:116); and AMB 1-2.15, AMB 2-1.15, AMB 4-3.15 and AMB 3-4.15 (SEQ ID NO:89, SEQ ID NO:101, SEQ ID NO:116 and SEQ ID NO:107).

Peptides of protein allergens of *Ambrosia artemisiifolia* within the scope of the invention can be used in methods of treating and preventing allergic reactions to ragweed pollen allergens. Thus, one aspect of the present invention provides therapeutic compositions comprising a peptide of *Amb a* I.1, *Amb a* I.2, *Amb a* I.3, *Amb a* I.4 or *Amb a* II including at least one T cell epitope, or preferably at least two T cell epitopes, and a pharmaceutically acceptable carrier or diluent. In another aspect, the therapeutic composition comprises a pharmaceutically acceptable carrier or diluent and a peptide comprising at least two regions, each region comprising at least one T cell epitope of a ragweed pollen allergen and is derived from the same or from different ragweed pollen allergens.

Administration of the therapeutic compositions of the present invention to desensitize an individual can be carried out using known techniques. For example, a peptide derived from a ragweed pollen allergen comprising at least one T cell epitope can be administered in combination with an appropriate diluent, or carrier. Preferably, peptides are administered in soluble form. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutically acceptable carriers include polyethylene glycol (Wie et al., *International Archives of Allergy and Applied Immunology* 64: 84-99 (1981)) and liposomes (Strejan et al., *Journal of Neuroimmunology* 7: 27 (1984)). For purposes of inducing T-cell non-responsiveness (or reduced T cell responsiveness) in an individual, the therapeutic composition is preferably administered in non-immunogenic form, e.g., one that does not

include adjuvant. Such compositions will generally be administered by injection (e.g., intravenous, subcutaneous, intramuscular), oral administration, (e.g., as in the form of a capsule), inhalation, transdermal application or rectal administration. Preferably, therapeutic compositions are administered subcutaneously.

5           In addition, isolated and purified native ragweed pollen protein allergens (see, Example 1 for isolation and purification of ragweed pollen protein antigens) or portions thereof, can be administered orally.

          The therapeutic compositions of the invention are administered to ragweed pollen-sensitive individuals at dosages and for lengths of time effective to reduce sensitivity (i.e.,  
10       reduce the allergic response) of an individual to a ragweed pollen allergen. A therapeutically effective amount of one or more of the same or of different therapeutic compositions can be administered simultaneously or sequentially to a ragweed pollen-sensitive individual. Effective amounts of the therapeutic compositions will vary according to factors such as the degree of sensitivity of the individual to ragweed pollen allergens, the age, sex, and weight of  
15       the individual, and the ability of the peptide to stimulate a T cell response in the individual.

          For subcutaneous injection of one or more therapeutic compositions of the invention, preferably about 1 mg- 3 mg and more preferably from about 20mg-1.5 mg, and even more preferably about 50 mg- 750 mg of each active component (peptide) per dosage unit may be administered. It is especially advantageous to formulate parenteral compositions in unit  
20       dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suited as unitary dosages for human subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the desired pharmaceutical carrier. The specification for the novel unit dosage forms of the invention are dictated by and directly  
25       dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of human subjects.

          Dosage regimen may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered over the course of days, weeks, months  
30       or years, or the dose may be proportionally increased or reduced with each subsequent injection as indicated by the exigencies of the therapeutic situation. In one preferred therapeutic regimen, subcutaneous injections of therapeutic compositions are given once a week for 3-6 weeks. The dosage may remain constant for each injection or may increase or decrease with each subsequent injection. A booster injection may be administered at  
35       intervals of about three months to about one year after initial treatment and may involve only a single injection or may involve another series of injections similar to that of the initial treatment.

          To administer a composition of the invention by other than parenteral administration, (i.e. oral administration) it may be necessary to coat the composition with, or co-administer

the composition with, a material to prevent its inactivation or enhance its absorption and bioavailability. For example, a peptide formulation may be co-administered with enzyme inhibitors or in liposomes. Enzyme inhibitors include pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEP) and trasylol. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al., (1984) *J. Neuroimmunol.*, 5 7:27). When a peptide is suitably protected, the peptide may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The peptide and other ingredients may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the individual's diet. For oral therapeutic administration, 10 the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, solutions, gels, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active compound. The percentage of the composition and preparations may, of course, be varied and may conveniently be between about 5 to 80% of the weight of the unit. The 15 amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. In addition, the active compound may be incorporated into sustained-release or controlled release (steady state or pulsatile release) preparations and formulations.

In yet another aspect of the present invention, a composition is provided comprising 20 at least two peptides (e.g., a physical mixture of at least two peptides), each comprising at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia*. Compositions comprising several peptides or combinations of separate peptides can include as many peptides as desired (e.g., 5, 6, 7...) for therapeutic efficacy. The peptides are derived from the same or from different ragweed pollen allergens. Such compositions can be administered in 25 the form of a therapeutic composition with a pharmaceutically acceptable carrier or diluent. Preferably, peptides are administered in soluble form. A therapeutically effective amount of one or more of such compositions can be administered simultaneously or sequentially, preferably subcutaneously, to a ragweed pollen-sensitive individual to desensitize or tolerize the individual to ragweed pollen.

30 Preferred compositions and preferred combinations of peptides which can be administered simultaneously or sequentially (comprising peptides having amino acid sequences shown in Fig. 14) include the following combinations: AMB 4-6.1 and RAE 70.1 (SEQ ID NO:111 and SEQ ID NO:44); AMB 4-6.1 and AMB 2-5.1 (SEQ ID NO:111 and SEQ ID NO:92); AMB 4-9.1 and AMB 2-5.1 (SEQ ID NO:110 and SEQ ID NO:92); AMB 4-9.1 and RAE 70.1 (SEQ. ID NO:110 and SEQ ID NO:44); AMB 4-6.1, AMB 2-5.1 and 35 AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:92 and SEQ ID NO:86); AMB 4-9.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:92 and SEQ ID NO:86); AMB 4-6.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:44 and SEQ ID NO:86); AMB 4-9.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:44 and SEQ ID NO:86); AMB 4-

6.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102); AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103); AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103); AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:101, and SEQ ID NO:116); AMB 1-2.15, AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:89, SEQ ID NO:101, and SEQ ID NO:116); and AMB 1-2.15, AMB 2-1.15, AMB 4-3.15 and AMB 3-4.15 (SEQ ID NO:89, SEQ ID NO:101, SEQ ID NO:116 and SEQ ID NO:107).

Particularly preferred compositions and preferred combinations of peptides for therapeutic administration include the following combinations: AMB 1-2.1 and AMB 4-9.1 (SEQ ID NO:86 and SEQ ID NO:110); AMB 1-2.1, AMB 2-38.1 and AMB 4-9.1 (SEQ ID NO:86, SEQ ID NO:141 and SEQ ID NO:110); AMB 1-2.1, AMB 2-38.1, AMB 4-9.1 and AMB 2-4.1 (SEQ ID NO:86, SEQ ID NO:141, SEQ ID NO:110 and SEQ ID NO:90); AMB 1-2.1, AMB 2-38.1, AMB 4-9.1, AMB 2-4.1 and AMB 3-5.1 (SEQ ID NO:86, SEQ ID NO:141, SEQ ID NO:110, SEQ ID NO:90, and SEQ ID NO:102); AMB 1-2.1, AMB 2-36.1 and AMB 4-9.1 (SEQ ID NO:86, SEQ ID NO:139 and SEQ ID NO:110); AMB 1-2.1, AMB 2-36.1, AMB 4-9.1 and AMB 2-4.1 (SEQ ID NO:86, SEQ ID NO:139, SEQ ID NO:110 and SEQ ID NO:90); and AMB 1-2.1, AMB 2-36.1, AMB 4-9.1, AMB 2-4.1 and AMB 3-5.1 (SEQ ID NO:86, SEQ ID NO:139, SEQ ID NO:110, SEQ ID NO:90, and SEQ ID NO:102).

Another aspect of this invention pertains to a multi-peptide formulation suitable for pharmaceutical administration to ragweed sensitive individuals. The multi-peptide formulation includes at least two or more peptides of ragweed pollen protein allergen having human T cell stimulating activity in an *in vitro* T cell proliferation assay (i.e., comprising at least one T cell epitope). Special considerations when preparing a multi-peptide formulation include maintaining the solubility and stability of all peptides in the formulation at a physiologically acceptable pH. This requires choosing one or more pharmaceutically acceptable carriers such as excipients which are compatible with all the peptides in the multi-peptide formulation. For example, suitable excipients include sterile water, sodium phosphate, mannitol or both sodium phosphate and mannitol or any combination thereof. Additionally due to the potential for dimerization of the peptides in a multi-peptide formulation, there may also be included an agent such as EDTA to prevent dimerization. Alternatively, any material or procedures known in the art to prevent dimerization may be

used. A preferred multi-peptide formulation includes at least one first peptide and at least one second peptide of ragweed pollen protein each having human T cell stimulating activity and soluble at a physiologically acceptable pH and selected from the group of peptides. In a preferred embodiment, the multi-peptide formulation includes Peptides Amb 1-2.1, Amb 4-9.1, Amb 2-36.1, and modifications thereof, and sodium phosphate and mannitol. In this embodiment, it is preferred that Peptides Amb 1-2.1, Amb 4-9.1, and Amb 2-36.1 are in the form of a lyophilized powder which is reconstituted in a physiologically acceptable carrier, such as sterile water, prior to use. As an illustrative example, a multi-peptide formulation comprising the three peptides were produced and used in Phase I human clinical trials (see Example XV). The peptides were combined during manufacturing to produce a vial containing a sterile, pyrogen free, lyophilized powder having the following composition:

<b>Active:</b>	Peptide Amb 1-2.1, Peptide Amb 4-9.1, and Peptide Amb 2-36.1 In concentration of 7.5-1500 µg per peptide
<b>Inactives:</b>	0.05 M Sodium Phosphate pH 7.5 5% w/v Mannitol, U.S.P.
<b>Diluent:</b>	Sterile Water for Injection, U.S.P. (initial reconstitution) 0.9% Sodium Chloride for Injection (dilution beyond initial reconstitution)

The multi-peptide formulation of the invention can be provided in the form of a kit, including instructions for use.

The present invention also provides methods of detecting sensitivity in individuals to ragweed pollen allergens comprising combining a blood sample obtained from the individual with a peptide of the present invention, under conditions appropriate for binding of blood components with the peptide and determining the extent to which such binding occurs. The extent to which binding occurs is determined by assessing T cell function, T cell proliferation or a combination thereof. Other diagnostic methods for allergic diseases which the protein or peptides of the invention can be used include radio-allergosorbent test (RAST), paper radioimmunosorbent test (PRIST), enzyme linked immunosorbent assay (ELISA), radioimmunoassays (RIA), immuno-radiometric assays (IRMA), luminescence immunoassays (LIA), histamine release assays and IgE immunoblots.

The presence in individuals of IgE specific for ragweed protein allergen and the ability of T cells of the individual to respond to T cell epitope(s) of the protein allergen can be determined by administering to the individuals an Immediate Type Hypersensitivity test and a Delayed Type Hypersensitivity test. The individuals are administered an Immediate Type Hypersensitivity test (see e.g. *Immunology* (1985) Roitt, I.M., Brostoff, J., Male, D.K. (eds), C.V. Mosby Co., Gower Medical Publishing, London, NY, pp. 19.2-19.18; pp. 22.1-22.10) utilizing purified native ragweed protein allergen, a peptide of ragweed protein, allergen or a modified form of the peptide, each of which binds IgE specific for the allergen. The same individuals are administered a Delayed Type Hypersensitivity test

prior to, simultaneously with, or subsequent to administration of the Immediate Type Hypersensitivity test. Of course, if the Immediate Type Hypersensitivity test is administered prior to the Delayed Type Hypersensitivity test, the Delayed Type Hypersensitivity test would only be given to those individuals exhibiting a specific Immediate Type Hypersensitivity reaction. The Delayed Type Hypersensitivity test utilizes a modified form of ragweed pollen protein or a portion thereof, ragweed protein allergen produced by recombinant DNA techniques, or peptide derived from ragweed pollen protein, each of which has the ability to stimulate human T cells and each of which does not bind IgE specific for the allergen in a substantial percentage of the population of individuals sensitive to the allergen (e.g., at least about 75%). After administration of the Delayed Type Hypersensitivity test, the extent to which a specific Delayed Type Hypersensitivity reaction occurs in the individual to the protein allergen or ragweed pollen protein peptide indicating presence in the individual of T cells specific to T cell epitope(s) of the protein allergen or ragweed pollen protein peptide is determined. Those individuals found to have both a specific Immediate Type Hypersensitivity reaction and a specific Delayed Type Hypersensitivity reaction are diagnosed as having sensitivity to a ragweed allergen and may, if need be, administered a therapeutically effective amount of a therapeutic composition comprising the modified form of ragweed protein allergen or portion thereof, the ragweed protein allergen produced by recombinant DNA techniques, or peptide, each as used in the Delayed Type Hypersensitivity test, and a pharmaceutically acceptable carrier or diluent.

It is also possible to modify the structure of a peptide of the invention for such purposes as increasing solubility, enhancing therapeutic or preventive efficacy, or stability (e.g., shelf life *ex vivo*, and resistance to proteolytic degradation *in vivo*.) A modified peptide can be produced in which the amino acid sequence has been altered, such as by amino acid substitution, deletion, or addition, to modify immunogenicity and/or reduce allergenicity, or to which a component has been added for the same purpose.

For example, a peptide can be modified so that it maintains the ability to induce T cell non-responsiveness and bind MHC proteins without the ability to induce a strong proliferative response or possibly, any proliferative response when administered in immunogenic form. In this instance, critical binding residues for the T cell receptor can be determined using known techniques (e.g., substitution of each residue such as, for example, with alanine and determination of the presence or absence of T cell reactivity). Those residues shown to be essential to interact with the T cell receptor can be modified by replacing the essential amino acid with another, preferably similar amino acid residue (a conservative substitution) whose presence is shown to enhance, diminish, but not eliminate, or not affect T cell reactivity. In addition, those amino acid residues which are not essential for T cell receptor interaction can be modified by being replaced by another amino acid whose incorporation may enhance, diminish or not affect T cell reactivity, but not eliminate

binding to relevant MHC. Preferred amino acid substitutions for non-essential amino acids include, but are not limited to substitutions with alanine, glutamic acid or a methyl amino acid.

Another example of a modification of peptides is substitution of cysteine residues preferably with serine, threonine, leucine or glutamic acid to minimize dimerization via disulfide linkages. As described in Example XI, a peptide from Region 2, RAE 70.1 (SEQ ID NO:44) was modified to minimize dimerization by substituting serine for the cysteine residue at amino acid position 212. In addition, as shown in Figure 23, peptide AMB 4-9.1 which contains an acid-sensitive aspartic acid-proline bond at amino acid residues 360-361, was modified to increase the stability of this peptide. For example, in peptide AMB 4-9.1DA, the proline at position 361 was substituted with alanine to remove the acid-sensitive bond.

Stability may also be enhanced in those peptides shown to be susceptible to degradation by deamidation (e.g. the peptide contains a labile Asn-Gly sequence susceptible to deamidation under various conditions. In such situations it has been found that lyophilization stabilized such peptides against deamidation. Peptide Amb 4-9.1 contained a labile Asn-Gly sequence and was found to be susceptible to deamidation under accelerated conditions (e.g. increased buffer concentration, increased ionic strength and increased temperature). Lyophilization stabilized the peptide against deamidation with no significant increase in degradants following 6 months of storage at 5°C.

In order to enhance stability and/or reactivity, peptides can also be modified to incorporate one or more polymorphisms in the amino acid sequence of a protein allergen resulting from natural allelic variation. Additionally, D-amino acids, non-natural amino acids or non-amino acid analogues can be substituted or added to produce a modified peptide within the scope of this invention. Furthermore, peptides can be modified using the polyethylene glycol (PEG) method of A. Schon and co-workers (Wie et al., *supra*) to produce a peptide conjugated with PEG. In addition, PEG can be added during chemical synthesis of a peptide of the invention. Modifications of peptides or portions thereof can also include reduction/alkylation (Tarr in: *Methods of Protein Microcharacterization*, J.E. Silver ed. Humana Press, Clifton, NJ, pp. 155-194 (1986)); acylation (Tarr, *supra*); esterification (Tarr, *supra*); chemical coupling to an appropriate carrier (Mishell and Shiigi, eds, *Selected Methods in Cellular Immunology*, WH Freeman, San Francisco, CA (1980); U.S. Patent 4,939,239); or mild formalin treatment (Marsh *International Archives of Allergy and Applied Immunology* 41: 199-215 (1971)).

In another embodiment, peptides within an allergen group (e.g., Amb a I or Amb a II) can be modified to enhance T cell reactivity. Given the cross-reactivity within the Amb a I family and Amb a II, a peptide of one group allergen which may be less reactive than a peptide of another group allergen corresponding in amino acid position can have one or more amino acids substituted with one or more amino acids from the corresponding peptide.

Additionally, peptides can be modified to incorporate a polymorphism in the amino acid sequence of a protein allergen resulting from natural allelic variation. Modification of peptides to include one or more of these polymorphisms may result in enhanced stability and/or reactivity.

5 To facilitate purification and potentially increase solubility of peptides of the invention, it is possible to add reporter group(s) to the peptide backbone. For example, poly-histidine can be added to a peptide to purify the peptide on immobilized metal ion affinity chromatography (Hochuli, E. et al., *Bio/Technology*, 6:1321-1325 (1988)). In addition, specific endoprotease cleavage sites can be introduced, if desired, between a reporter group  
10 and amino acid sequences of a peptide to facilitate isolation of peptides free of irrelevant sequences. In order to successfully desensitize an individual to a protein antigen, it may be necessary to increase the solubility of a peptide by adding functional groups to the peptide or by not including hydrophobic T cell epitopes or regions containing hydrophobic epitopes in the peptides. For example, in Region 3 of the *Amb a* 1.1 protein, a selected peptide AMB 3-  
15 4.1 was modified to increase its solubility by the addition of three naturally occurring sequential residues found in the *Amb a* 1.1 protein, "RHG", to the 5' end of the peptide. These residues are not necessary for T cell recognition and are also found in peptide AMB 3-5.1. Similarly, as described in Example XI, peptide RAE 70.1 was divided into two fragments and modified to increase solubility. As shown in Figure 25, a fragment of RAE  
20 70.1 ("ORIGINAL" in Figure 25, amino acid residues 194-216, SEQ ID NO:125) was modified by substitution or addition of charged amino acids to increase the hydrophilicity and decrease the pI of the peptide to thereby increase the solubility. For example, in peptide Amb 2-22.1, isoleucine was substituted with glutamic acid to decrease the pI and avoid precipitation of the peptide from solution at a physiological pH. Similar substitutions and  
25 additions are shown in Figure 25. In addition, charged amino acids or charged amino acid pairs or triplets when added to the carboxy or amino terminus of the peptide may be particularly useful to increase the solubility of the peptide. Examples of charged amino acids include, but are not limited to arginine (R), lysine (K), histidine (H), glutamic acid (E), and aspartic acid (D). Examples of such modifications are shown in Fig. 14, AMB 2-8.1, AMB  
30 2-9.1, AMB 2-10.1, AMB 2-7.1 and AMB 2-11.1.

To potentially aid proper antigen processing of T cell epitopes within a peptide, canonical protease sensitive sites can be recombinantly or synthetically engineered between regions, each comprising at least one T cell epitope. The resulting peptide can be rendered sensitive to cathepsin and/or other trypsin-like enzymes cleavage to generate portions of the  
35 peptide containing one or more T cell epitopes. In addition, such charged amino acid residues can result in an increase in solubility of a peptide.

Site-directed mutagenesis of DNA encoding a peptide of the invention can be used to modify the structure of the peptide. Such methods may involve PCR with degenerate oligonucleotides (Ho et al., *Gene*, 77:51-59 (1989)) or total synthesis of mutated genes



(Hostomsky, Z., et al., *Biochem. Biophys. Res. Comm.*, 161:1056-1063 (1989)). To enhance bacterial expression, the aforementioned methods can be used in conjunction with other procedures to change the eucaryotic codons in DNA constructs encoding peptides of the invention to ones preferentially used in *E. coli*, yeast, mammalian cells or other eucaryotic cells.

Another aspect of the invention pertains to an antibody specifically reactive with *Amb a I*, or a fragment thereof. The antibodies of this invention can be used to standardize allergen extracts or to isolate the naturally-occurring or native form of *Amb a I*. For example, by using proteins or fragments thereof based on the cDNA sequence of *Amb a I*, anti-protein/anti-peptide antisera or monoclonal antibodies can be made using standard methods. A mammal such as a mouse, a hamster or rabbit can be immunized with an immunogenic form of such protein or an antigenic fragment which is capable of eliciting an antibody response. Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques well known in the art. *Amb a I* or fragment thereof can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as antigen to assess the levels of antibodies.

Following immunization, anti-*Amb a I* antisera can be obtained and, if desired, polyclonal anti-*Amb a I* antibodies isolated from the serum. To produce monoclonal antibodies, antibody-producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to yield hybridoma cells. Such techniques are well known in the art, for example the hybridoma technique originally developed by Kohler and Milstein, (*Nature* (1975) 256:495-497) as well as other techniques such as the human B cell hybridoma technique (Kozbar et al., *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy* (1985) Alan R. Liss, Inc. pp. 77-96). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with *Amb a I* and the monoclonal antibodies isolated.

The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with *Amb a I*. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')<sub>2</sub> fragments can be generated by treating antibody with pepsin. The resulting F(ab')<sub>2</sub> fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The antibody of the present invention is further intended to include bispecific and chimeric molecules having an anti-*Amb a I* portion.

Another aspect of this invention provides T cell clones and soluble T cell receptors specifically reactive with *Amb a I* or a fragment thereof. Monoclonal T cell populations (i.e., T cells genetically identical to one another and expressing identical T cell receptors) can be

derived from an individual sensitive to *Amb a I*, followed by repetitive *in vitro* stimulation with *Amb a I* or portion thereof in the presence of MHC-matched antigen-presenting cells. Single *Amb a I* MHC responsive cells can then be cloned by limiting dilution and permanent lines expanded and maintained by periodic *in vitro* restimulation. Alternatively, *Amb a I* specific T-T hybridomas can be produced by a technique similar to B cell hybridoma production. For example, a mammal, such as a mouse can be immunized with *Amb a I* or fragment thereof, T cells from the mammal can be purified and fused with an autonomously growing T cell tumor line. From the resulting hybridomas, cells responding to *Amb a I* or fragment thereof are selected and cloned. Procedures for propagating monoclonal T cell populations are described in *Cellular and Molecular Immunology* (Abul K. Abbas et al. ed.), W.B. Saunders Company, Philadelphia, PA (1991) page 139. Soluble T cell receptors specifically reactive with *Amb a I* or fragment thereof can be obtained by immunoprecipitation using an antibody against the T cell receptor as described in *Immunology: A Synthesis* (Second Edition), Edward S. Golub et al., ed., Sinauer Associates, Inc., Sunderland, MA (1991) pages 366-269.

T cell clones specifically reactive with *Amb a I* or fragment thereof can be used to isolate and molecularly clone the gene encoding the relevant T cell receptor. In addition, a soluble T cell receptor specifically reactive with *Amb a I* or fragment thereof can be used to interfere with or inhibit antigen-dependent activation of the relevant T cell subpopulation, for example, by administration to an individual sensitive to a cat allergen. Antibodies specifically reactive with such a T cell receptor can be produced according to the techniques described herein. Such antibodies can be used to block or interfere with the T cell interaction with peptides presented by MHC.

The present invention also provides nucleic acids having sequences encoding proteins and peptides of the invention. Nucleic acid sequences used in any embodiment of this invention can be cDNA as described herein, or alternatively, can be any oligodeoxynucleotide sequence having all or a portion of a sequence represented herein, or their functional equivalents. Such oligodeoxynucleotide sequences can be produced chemically or mechanically, using known techniques. A functional equivalent of an oligonucleotide sequence is one which is 1) a sequence capable of hybridizing to a complementary oligonucleotide to which the sequence (or corresponding sequence portions) of SEQ ID NO: 1, 3, 5, 7, 9 and 11 or fragments thereof hybridizes, or 2) the sequence (or corresponding sequence portion) complementary SEQ ID NO: 1, 3, 5, 7, 9 and 11 and/or 3) a sequence which encodes a product (e.g., a polypeptide or peptide) having the same functional characteristics of the product encoded by the sequence (or corresponding sequence portion) of SEQ ID NO: 1, 3, 5, 7, 9 and 11. Whether a functional equivalent must meet one or more criteria will depend on its use (e.g., if it is to be used only as an oligoprobe, it need meet only the first or second criteria and if it is to be used to produce a peptide of the present invention, it need only meet the third criterion).

As described in the Examples which follow, *Amb a I* and *Amb a II* proteins have been recombinantly expressed in *E. coli*, purified and shown to have reduced binding to human allergic ragweed pollen IgE on Western blots. Overlapping peptides derived from the *Amb a I.1* protein and various peptides derived from *Amb a I.3* and *Amb a I.2* were synthesized and used to identify regions of T cell reactivity within the protein. These regions of T cell reactivity were further defined by modifying selected *Amb a I.1* peptides and determining T cell reactivity to these peptides.

This invention is further illustrated by the following non-limiting examples.

#### 10 **Example I Native Ragweed Pollen Allergen Purification**

What follows is a description of the work done to biochemically purify the allergens of *Ambrosia artemisiifolia* in their native form as primary antigens for human T-cell epitope mapping:

50 g of defatted short ragweed pollen (Greer Labs) was extracted in 500 ml .05 M Tris pH 7.95 containing protease inhibitors. The extract was then depigmented by batch absorption with Whatman DE-52 DEAE cellulose (150 g dry weight) in the presence of 0.2 M NaCl at 4°C. Unbound material was dialysed against .025 M Tris pH 7.95 with protease inhibitors. The depigmented sample was next applied to an 80 ml DEAE cellulose column (Whatman DE-52) equilibrated in .025 M Tris pH 7.95 containing protease inhibitors. Acidic proteins were eluted with .025 M Tris, 0.2 M NaCl pH 7.95 at 4°C with inhibitors.

In order to biochemically purify *Amb a I*, the acidic DEAE elution sample was fractionated by ammonium sulfate precipitation into 0-45% and 45-59% saturation samples (4°C). The *Amb a I*-enriched 45-59% pellet was applied at 0.5 ml/min (4°C) to a 500 ml Sephacryl S200 (Pharmacia) column in .05 M ammonium bicarbonate containing inhibitors. Purified *Amb a I* was recovered in the 38 kD region and dialysed against .04 M Tris pH 8.0. This sample was next applied to an 8 ml Mono Q HR 10/10 (Pharmacia) column in .04 M Tris pH 8.0 at 25°C. Elution was performed with .04 M Tris pH 8.0 containing .08M NaCl and the major peaks were analyzed as discussed below for confirmation and purity of *Amb a I*.

To biochemically purify *Amb a II*, the acidic DEAE elution sample was separated into an *Amb a II*-enriched fraction by ammonium sulfate precipitation at 0-45% saturation (4°C). The pellet was applied at 0.5 ml/min (4°C) to a 200 ml Sephadex G75 (Pharmacia) column in .05 M ammonium bicarbonate containing inhibitors. Purified *Amb a II* was recovered in the 38 kD region and dialysed against .04 M Tris pH 8.0 at 25°C to separate contaminating *Amb a I* from *Amb a II*. Elution was performed with .04 M Tris pH 8.0 containing .08M NaCl.

The major peaks were analyzed by IEF SDS-PAGE using a mixture of 4.5-5.3 and 3.5-10.0 ampholytes (Pharmacia) in a 7% acrylamide gel. Protein sequencing was also performed to confirm *Amb a II*.

In another embodiment *Amb a* I.1 and *Amb a* I.2 were biochemically purified as follows. Defatted short ragweed pollen was purchased from Greer laboratories (Lenior, North Carolina). The pollen (50g) was extracted overnight at 4°C with extraction buffer (10ml/g pollen) containing 50 mM Tris-HCl, pH8 and protease inhibitors in final concentrations: phenyl methyl sulfonyl fluoride (170 µg/ml); soybean trypsin inhibitor (1 µg/ml); leupeptin (1 µg/ml) and pepstatin A (1 µg/ml). The soluble extract was clarified by sequential filtration through Whatman #1 paper (Whatman, Maidstone England) followed by an 0.8 micron filter. The soluble pollen extract (SPE) was either used for IgE binding studies or subjected to further purification as described below. For purification of *Amb a* I.1 and *Amb a* I.2, sodium chloride was added to SPE to a final concentration of 0.3M and the material was depigmented by batch absorption onto What DE 52 cellulose (50 g dry weight) with extraction buffer plus 0.3 M NaCl. The solution was separated from the resin by filtration (Whatman #1 paper, Whatman, Maidstone England). The sample was then fractionated by sequential (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> precipitation, first at 0-47.5% and then at 47.5%-60% saturation at 4°C. The latter precipitate enriched in *Amb a* I, was separated by centrifugation (10,000 x g, 1h) and the pellet was resuspended in 20 mM Tris-HCl, pH 8. The sample was then loaded onto an anti-*Amb a* I murine monoclonal antibody affinity column (mAb-4B5B7, Dr. D. Klapper, Univ. North Carolina, Chapel Hill, NC) which was determined to bind members of the *Amb a* I/*Amb a* II family. *Amb a* I was eluted with 0.1 M glycine, pH11 and was immediately neutralized with 1 M sodium phosphate, pH 3.9. This preparation contained both *Amb a* I.1 and *Amb a* I.2 as detected by SDS-PAGE analysis and by Western blotting. The affinity purified proteins were pooled, concentrated and dialyzed against 20 mM Tris, pH 8.5, extensively. *Amb a* I.1 and *Amb a* I.2 were further separated by a Source Q ionic exchange column (Pharmacia, Piscataway, NJ), which was equilibrated with 20 mM Tris, pH 8.5. The flow-through material contained *Amb a* I.2 whereas *Amb a* I.1 was eluted with step gradients of 24 mM and 80 mM NaCl. The latter peak also contained some α,β cleaved fragments of *Amb a* I.1. *Amb a* I.1 and *Amb a* I.2 purity was verified by NH<sub>2</sub>-terminal amino acid sequencing as being greater than 95%.

### 30 **Example II Recombinant Ragweed Pollen Allergen Expression**

What follows is a description of the work done to produce the allergens of *Ambrosia artemisiifolia* as recombinant proteins in *E. coli*.

Described and provided in USSN 07/529,951, filed May 29, 1990, (incorporated herein by reference) are full length cDNAs encoding *Amb a* I.1, *Amb a* I.2, *Amb a* I.3, *Amb a* I.4 and *Amb a* II. cDNA inserts encoding these five protein allergens were constructed in the vector pTrc99A (Amann, E. et al., *Gene*, 69:301 (1988)) which was kindly provided by Dr. Egon Amann (Behringwerke AG, Marburg, FRG). The nucleotide sequences and deduced amino acid sequences of the *Amb a* I family members are shown in the sequence listing as SEQ ID NO:1 and 2 (*Amb a* I.1), SEQ ID NO:3 and 4 (*Amb a* I.2), SEQ ID NO:5 and 6

(*Amb a* I.3), SEQ ID NO:7 and 8 (*Amb a* I.4), and SEQ ID NO:9 and 10 (*Amb a* II). The cDNAs encoding each allergen were cloned in frame with a polylinker encoding six sequential histidines, (CAC)<sub>6</sub>, that had been inserted into the 5' end of the pTrc99A vector as a NcoI/EcoRI synthetic adapter (Maniatis T., et al.

5 *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982). These cDNA inserts were then cloned in-frame into the appropriate pTrc99. To further enhance expression, a retroregular stem-loop sequence was placed at the 3' end in the untranslated region (Skoglund, C. M., et al. *Gene* 88:1 (1990)). The H<sub>6</sub> leader sequence allowed purification using QIAGEN NTA-Agarose (Diagen GmbH, Dusseldorf, FRG), a Ni<sup>2+</sup> chelating support (Hochuli, E. et al., *Biotechnology*, 6:1321 (1988)). The  
10 vectors were transformed into the XL-1 Blue host bacteria. Expression of individual recombinant proteins was induced when cultures reached OD<sub>600</sub>=0.6 to 0.7 by the addition of isopropyl-D-thiogalactopyranoside (IPTG) to 1 mM concentration to the culture medium. After 2 hours of further growth at 37°, the cells were pelleted. Recombinant proteins were  
15 then obtained by either of the following procedures. In one embodiment, the cells were resuspended in lysozyme containing phosphate buffer (0.4 mg/ml) and incubated for 30 minutes on ice. The cell suspension was frozen and quick thawed followed by sonication (Bond, J. G. et al., *J. Immunol.*, 146:3380 (1991)). Insoluble recombinant protein was recovered by a low speed centrifugation and solubilized in 10 ml (per 1 liter growth) of  
20 buffer containing 8 M urea, 50 mM Tris, pH 8.0, 2 mg/ml leupeptin, 2 mg/ml pepstatin and 1 mg/ml soybean trypsin inhibitor. The urea solubilized preparation was subjected to a low speed centrifugation and the recombinant proteins in the supernatant isolated by metal ion chromatography (Hochuli *supra*).

In another embodiment, the pelleted bacteria were resuspended in 6 M guanidine  
25 HCl, 100 mM 2-mercaptoethanol, 100 mM NaPO<sub>4</sub>, 10 mM Tris pH 8.0. This suspension was subjected to centrifugation at 15,000 X g, and the supernatant removed, adjusted to pH 8.0 with 10 N NaOH, and applied to an NTA agarose column that had been equilibrated in 6 M guanidine HCl, 100 mM NaPO<sub>4</sub>, 10 mM Tris pH 8.0. The column was washed in 6 M  
30 guanidine HCl, 100 mM NaPO<sub>4</sub>, 10 mM Tris pH 8.0 until the OD<sub>280</sub> of the effluent reached background. The column buffer was then switched to 8 M urea, 100 mM NaPO<sub>4</sub>, 10 mM Tris pH 8.0. After equilibration, a more stringent wash was performed in 8 M urea, 100 mM NaOAc, 10 mM Tris pH 6.3 until the OD<sub>280</sub> of the effluent reached background. Recombinant protein (as an H<sub>6</sub> fusion) was then eluted in 8 M urea, 100 mM NaOAc, 10  
35 mM Tris pH 4.5 and collected in aliquots whose OD<sub>280</sub> profile was monitored. The protein peak was dialyzed 3 times into 500 volumes of PBS for human T-cells analysis. Yield ranged from 1-3 mg of recombinant protein per litre with purity of approximately 55% (as determined by densitometric scanning).

In another embodiment, high yields (e.g. 50-100 mg purified r *Amb a* I.1 protein/L of fermentation broth) were recovered. This embodiment involves the use of a construct which

is different from the one described above in 3 ways: 1) deletion of sequence coding for 36 non-*Amb a I* related amino acids, 2) attachment of a 6 histidine linker to the NH<sub>2</sub>-terminus, and 3) replacement of 2 existing arginine codons with *E. coli*-preferred arginine codon. The resulting construct showed significantly higher expression levels than did the original construct. This phenomenon was true in both shaker flask and 10L fermentation cultures. The purification also differed significantly from the procedure described above. Cell paste from fermentors was homogenized and inclusion bodies were purified prior to solubilization in guanidine-HCL. The solubilized inclusion bodies were then purified using metal chelate chromatography and buffer exchanged into acetate buffer. The resulting protein is greater than 90% full length r *Amb a I.1*.

### **Example III IgE Analysis With Purified Recombinant Ragweed Pollen Allergens**

Both Western blotting and ELISA techniques were used to analyze the binding of IgE to purified recombinant *Amb a I* and *Amb a II* proteins. Additionally, ELISA was used for analysis of IgE reactivity to peptides derived from the *Amb a I.1* protein.

#### **A. Western Blot Analysis**

The antigens were loaded on the gels as follows: lane 1 SPE (Soluble Pollen Extract), 15 mg/lane; lane 2 r*Amb a I.1* (recombinant *Amb a I.1*), 3 mg/lane; lane 3 r*Amb a I.2*, 4 mg/lane; lane 4 r*Amb a I.3*, 3 mg/lane; lane 5 r*Amb a I.4*, 3 mg/lane; and lane 6 r*Amb a II.1*, 4 mg/lane.

The gel electrophoresis and Western blot transfer procedures used were essentially as described elsewhere (Towbin, H., et al., *Proc. Natl. Acad. Sci., USA*, 76:4350 (1979)). Briefly, SDS-PAGE was performed in 10% acrylamide gels under reducing conditions (10mM dithiothreitol, at constant current). The transfer to nitrocellulose (0.1 mM, Schleicher & Schuell, Keene, NH) was performed in a Hoeffer apparatus according to the protocol of Towbin et al., *supra*. After transfer, the blots were rinsed in blot solution (25 mM Tris-HCl, pH 7.5, 0.17 M NaCl, and 0.05% Tween 20) and stained for 1 hour with 0.1% India ink. All subsequent incubations with antibodies and washes were performed in blot solution at room temperature. The first antibody incubations were performed overnight then rinsed and incubated with the appropriate biotinylated second antibody (Kirkegaard Perry Laboratories, Gaithersburg, MD). The final incubation was performed using <sup>125</sup>I streptavidin (1 mCi/25 ml blot solution) for 1 hour followed by removal of unbound labeled material and autoradiography at -80°C with an intensifying screen.

Fig. 1 shows the Western blot IgE binding pattern of two ragweed allergic patient plasma samples (#475 and #143). These patterns are representative of the typical IgE reactivity to these proteins on a Western blot. Both patients show binding to the *Amb a I.1* and *Amb a I.3* gene products in the soluble pollen extract (lane 1, Fig. 1). The results

demonstrate clear IgE binding reactivity in these patients to *Amb a* I.1, *Amb a* I.3, *Amb a* I.4. In contrast, *Amb a* I.2 and *Amb a* II.1 showed markedly reduced IgE binding.

### B. ELISA

5 Patient #143 IgE was also tested for binding to purified recombinant *Amb a* I and *Amb a* II proteins by ELISA. Fig. 2 shows the results of this analysis performed according to the following method.

Corning assay plates (#25882-96) were coated with each antigen listed in Fig. 2 at the following concentrations: Soluble Pollen Extract (SPE) 15 mg/ml; *rAmb a* I.1, 5 mg/ml; 10 *rAmb a* I.2, 20 mg/ml; *rAmb a* I.3, 5 mg/ml; *rAmb a* I.4, 15 mg/ml; and *rAmb a* II, 20 mg/ml. 50 mls/well of the above antigens were added and coating was carried out overnight at 4° C. The coating antigens were removed and the wells were blocked with 0.5% gelatin in PBS, 200ml/well for 2 hours at room temperature. Patient #143 plasma was serially diluted with PBS-Tween 20 (PBS with 0.05% nonionic detergent Tween-20 Sigma, St. Louis MO) and 15 100ml/well was added and incubated overnight at 4°C (plasma dilutions are tested in duplicate). The second antibody (biotinylated goat anti-Human IgE, 1:1000, Kirkegaard & Perry Laboratories Inc. Gaithersburg, MD), was added at 100 ml/well for one hour at room temperature. This solution was removed and streptavidin-HRPO, 1:10000, (Southern Biotechnology Associates, Inc., Birmingham, AL) was then added at 100ml/well for one 20 hour at room temperature (all wells are washed three times with PBS-Tween between each incubation step). TMB Membrane Peroxidates Substrate system (Kirkegaard & Perry Laboratories) was freshly mixed, and added at 100ml/well. The color was allowed to develop for 2-5 minutes. The reaction was stopped by the addition of 100ml/well of 1 M phosphoric acid. Plates were read on a Microplate EL 310 Autoreader (Biotek Instruments, 25 Winooski, VT) with a 450nm filter. The absorbance levels of duplicate wells was averaged. The graphed results (log of the dilution vs absorbance) of the ELISA assays are shown in Fig. 2. The order of coating antigens listed vertically in these figures corresponds in order from left to right to the coating antigens listed for each histogram.

The results of the ELISA assay, Fig. 2, demonstrate a similar reactivity pattern as 30 seen in the Western blot analysis (Fig. 1). The observation of much stronger binding to SPE is due to the fact that this antigen preparation is non-denatured. However, the recombinant *Amb a* proteins, because of their non-native bacterial origin behave as denatured antigens. The hierarchy of IgE binding in this patient to *rAmb a* proteins by ELISA shows that *Amb a* I.1 binds IgE more strongly than *Amb a* I.3 and *Amb a* I.4. The results also show that *Amb a* I.2 and *Amb a* II demonstrate markedly reduced binding to IgE. 35

As shown in Fig. 3, another ELISA assay was performed using a pool of human allergic sera. This assay demonstrates that biochemically purified (native) *Amb a* I and *Amb a* II proteins bind significantly more IgE than the respective recombinant proteins (*rAmb a* I.1 and *rAmb a* II). (The control in this experiment was plates coated without antigen and PBS).

### C. Modified ELISA for Greater Sensitivity

In another embodiment, an ELISA based assay system for serum IgE binding to peptides was developed which had greater sensitivity than direct ELISA, and creating tandem  
 5 *Amb a* I peptide copies in the form of a recombinant trimer. One strategy for constructing a recombinant trimer based on *Amb a* I peptide Amb 4-9.1 is as follows.

DNA sequences encoding Amb 4-9.1 are amplified by PCR from a plasmid containing *Amb a* I.1 sequence. Three Amb 4-9.1 copies are made each containing unique  
 10 restriction enzyme cleavage sites. The restriction sites are designed (inserted during the amplification reaction) to ensure peptide ligation to each other or ligation to plasmid. Amb 4-9.1 copies are ligated together and then the fused sequences are ligated into a sequencing vector, pUC19. After DNA sequence confirmation, the trimer sequences are excised from pUC19 and inserted into an expression vector pEt-11d containing a 6 histidine tag.

After DNA sequence confirmation, the expression plasmid containing Amb 4-9.1  
 15 trimer sequence was expressed in *E. coli* BL21. Expressed protein was purified over a  $\text{Ni}^{2+}$  column, protein was analyzed by SDS-Page, Western, ELISA and protein sequencing. The Amb 4-9.1 trimer was greater than 95 % pure and was soluble in PBS. The sequence of the Amb 4-9.1 trimer was confirmed as:

MGHHHHHHEFELGTKDVLNGAIFVASGVDPVLTPEQSAGSRGTK  
 20 DVLNGAIFVASGVDPVLTPEQSAGLQGTKDVLNGAIFVASGVDPVLTPEQSAG.

Side by side direct ELISAs were carried out in order to compare the sensitivity of assays using unmodified Amb 4-9.1 peptide versus recombinant Amb 4-9.1 trimer  
 constructed as discussed above. These assays were carried out using a panel of plasma samples from individuals with high skin test reactivity to *Amb a* I. Plasma previously shown  
 25 to have IgE binding to unmodified Amb 4-9.1 peptide served as a positive control. When IgE binding the the Amb 4-9.1 recombinant trimer was detected, competition ELISA was used to verify that the binding was relevant to the unmodified Amb 4-9.1 peptide (rather than to a novel epitope created by the recombinant Amb 4-9.1 structure).

There was a significant enhancement of assay sensitivity (both the number of  
 30 positives and assay signal/noise ratio) using the Amb 4-9.1 recombinant trimer. Thus, a very sensitive assay has been developed which is useful for screening IgE binding to any *Amb a* pollen protein allergen peptide. This assay is particularly useful for screening *Amb a* I.1 peptides which are to be formulated into a human therapeutic. This modified assay is also useful for screeninf for IgE binding to any peptide derived from any antigen by first  
 35 constructing a recombinant trimer of the peptide of interest as discussed above and using the recombinant trimer in the assay for detecting IgE binding.



**Example IV Tolerization of Mice with Recombinant *Amb a* I Proteins**

Balb/c mice (H-2 d) were immunized in the hind foot pads and at the base of the tail with an emulsion of Complete Freund's Adjuvant (CFA) and 50mg/mouse *Amb a* I.1. Seven days later, the popliteal lymph nodes, superficial inguinal lymph nodes, and the periaortic lymph nodes were harvested and cultured (1 x 10<sup>5</sup> lymph node cells + 2 x 10<sup>5</sup> irradiated spleen feeders) with various challenge antigens *in vitro*. The *in vitro* antigens consisted of various doses of *Amb a* I.1, *Amb a* I.2, *Amb a* I.3, *Amb a* I.4, *Amb a* II, and a media control. Cultures were incubated for 3 days at 37°C in a CO<sub>2</sub> incubator. On day 3, each culture was pulsed with 1mCi <sup>3</sup>H-thymidine and on day 4 the cultures were harvested and proliferation was monitored by incorporation of <sup>3</sup>H into DNA.

As seen in Fig. 4A, the mice that were immunized with 50mg *Amb a* I.1 + CFA have a good response to *Amb a* I.1, *Amb a* I.3, and *Amb a* I.4. The response to *Amb a* I.2 and *Amb a* II is also good but less than other antigens. It appears as though *Amb a* I.1 is very immunogenic in Balb/c mice and results in a response to each *Amb a* family member. When mice were immunized with PBS + CFA, there was no significant response to any of the *Amb a* family members (Fig. 4B).

An experiment was conducted to determine whether mice could be tolerized with *Amb a* I.1. The mice were tolerized with *Amb a* I.1 and pollen extract and then challenged with *Amb a* I.1 or pollen extract. The outline of this experiment #5 is shown below:

- Day 1:           Group (1): 6 Balb/c mice tolerized with 300mg *Amb a* I.1 + Incomplete Freund's Adjuvant (IFA) intraperitoneally (i.p.)  
                  Group (2): 6 Balb/c mice tolerized with 300mg Pollen Extract + IFA i.p.  
                  Group (3): 6 Balb/c mice exposed to PBS + IFA i.p.
- Day 15           Challenge 3 mice from each group with 50mg *Amb a* I.1 + CFA subcutaneously (s.c.)  
                  Challenge the remaining 3 mice from each group with 50mg Pollen extract + CFA s.c.
- Day 23           Harvest Lymph Nodes (Analyze each mouse separately)  
                  Test Lymph Node Response on *Amb a* I.1, I.2, I.3, I.4, II.1, Pollen Extract, Concanavalin A (Con A), and media.

The animals were sacrificed by cervical dislocation on day 23 and the popliteal lymph nodes, superficial inguinal lymph nodes, and the periaortic lymph nodes were removed and placed in rinsing buffer (cold RPMI 1640 containing 1% FCS). The nodes were rinsed with rinsing buffer and forced through a fine stainless steel mesh, using a glass pestal to suspend them in rinsing buffer. The suspended cells were rinsed two times by centrifugation at 1000

- 32 -

rpm for 10 minutes and resuspended in rinsing buffer. An aliquot from each sample was taken in order to do a cell count. The cells ( $4 \times 10^6$ /ml) were incubated in culture media (RPMI 1640 media containing 10% FCS, 2mM L-glutamine, 50U/ml penicillin, 50 mg/ml streptomycin and  $5 \times 10^{-5}$  M 2-mercaptoethanol) and test antigens at various concentrations.

5 The triplicate 0.1 ml cultures (U-bottom 96 well plates (Costar)) were incubated at 37°C and at 5% CO<sub>2</sub>. After 24 hours, 50ml of media from each well was placed in separate flat bottom 96 well plates (Costar) and frozen overnight at -20°C to eliminate carryover of live cells.

10 The supernatants were tested after thawing for their ability to support the growth of CTLL3, an IL-2 dependent T cell clone. CTLL3 in log phase growth were rinsed 3 times by centrifugation at 1000 rpm for 10 minutes. CTLL3's were added to the warmed culture supernatants ( $5 \times 10^3$  cells/well) and the IL-2 assay was incubated at 37°C and 5% CO<sub>2</sub>. After 24 hours, 1 mCi/well <sup>3</sup>H thymidine was added in 50 ml/well and the CTLL3 cells were incubated for an additional 4-6 hours. Following the pulse with <sup>3</sup>H, the plates were harvested on a Tom Tek 96 well cell harvester. The <sup>3</sup>H incorporation in each well was

15 counted by a Betaplate Model 1205 scintillation counter. Background counts were not subtracted.

Fig. 5 shows the lymph node responses to *Amb a* I.1, *Amb a* I.2, *Amb a* I.3, *Amb a* I.4, and *Amb a* II in the mice that were tolerized with *Amb a* I.1 or Phosphate Buffered Saline (PBS) and challenged with *Amb a* I.1. Fig. 5A, shows that the mice that were tolerized with

20 *Amb a* I.1 have a lower response to *Amb a* I.1 than the mice that were tolerized with PBS. The same appears to be true in the response to *Amb a* I.2, *Amb a* I.3, and *Amb a* I.4 (Figs. 5B, 5C, 5D). In Figs. 5E and 5F, it appears as though there is no significant difference in the response to *Amb a* II or pollen extract between the tolerized group and the non-tolerized group. Figs. 6A-6F show the lymph node responses to the various proteins in the mice that

25 were exposed to pollen extract or PBS and challenged with pollen extract. Table 1 summarizes the results of these experiments.

TABLE 1

	<u>Amba I.1</u>	<u>Amba I.2</u>	<u>Amba I.3</u>	<u>Amba I.4</u>	<u>Amba II.1</u>	<u>Pollen</u>
30 Tolerize/Challenge						
Control/ <u>Amba I.1</u>	+	+	+	+	-	-
<u>Amba I.1</u> / <u>Amba I.1</u>	-	-	-	-	-	-
35 Control/Pollen	+	+	+/-	+/-	-	+
Pollen/Pollen	-	-	-	-	-	-

(-) indicates a diminished response as compared to control

**Example V Synthesis of Overlapping Peptides**

*Amb a* I.1 overlapping peptides as shown in Fig. 7 were synthesized using standard Fmoc/tBoc synthetic chemistry and purified by dialysis or Reverse Phase HPLC. In addition, various peptides derived from *Amb a* I.2 and *Amb a* I.3 were synthesized. The amino acid residues of the synthesized peptides are in brackets by the peptide name and the amino acid sequence (in single letter code) is next to the peptide name. The peptide names are consistent throughout the Figures. In the design of the overlapping peptides, the relationship of the *Amb a* I family members at the level of T cell cross-reactivity as determined in Example VIII was considered. As shown in Table IV, the *Amb a* I protein allergens share a high degree of cross-reactivity. In addition, *Amb a* I.1 and *Amb a* II were found to have 55.2% cross-reactivity. Given the high degree of cross-reactivity it was expected that "shared" T cell epitopes exist among Groups I and II. Thus, the amino acid sequences of the *Amb a* I family members and *Amb a* II were examined to identify conserved and variable regions. It was expected that conserved regions within the *Amb a* I family members and *Amb a* II would contain "shared" T cell epitopes.

**Example VI T Cell Responses to Ragweed Peptides**

Peripheral blood mononuclear cells (PBMC) were purified by Sepacell-MN or lymphocyte separation medium (LSM) centrifugation of 60 ml of heparinized blood from ragweed-allergic patients who exhibited clinical symptoms of seasonal rhinitis and were skin prick test positive for ragweed. T cell lines were established from these cells by stimulation of  $1-2 \times 10^6$  PBMC/ml in RPMI-1640 containing 5% human AB serum (complete medium) with recombinant *Amb a* I.1 at 20-30 mg/ml for 5-6 days at 37°C in a humidified CO<sub>2</sub> incubator. Viable cells were purified by LSM centrifugation and cultured in complete medium supplemented with 5 units recombinant human IL-2/ml and 5 units recombinant human IL-4/ml for up to three weeks until the cells no longer responded to lymphokines and were considered "rested". The ability of the T cells to proliferate to *Amb a* I.1, *Amb a* I.2 and *Amb a* I.3 sequence-derived synthetic peptides was then assessed.

For assay,  $2 \times 10^4$  rested cells were restimulated in the presence of  $2 \times 10^4$  autologous Epstein-Barr virus (EBV)-transformed B cells (gamma-irradiated with 25,000 RADS) or  $5 \times 10^4$  autologous PBMC (3,500 RADS) with various concentrations of *Amb a* I.1 synthetic peptides in a volume of 200 µl complete medium in duplicate or triplicate wells in 96-well round bottom plates for 3 days. Each well then received 1 mCi tritiated thymidine for 16-20 hours. The counts incorporated were collected onto glass fiber filter mats and processed for liquid scintillation counting. Table II shows the results of a representative assay. The maximum response in a titration of each peptide is expressed as the S.I. or stimulation index. The S.I. is the CPM incorporated by cells in response to peptide divided by the CPM incorporated by cells in medium only. An S.I. value greater than the background level is considered "positive" and indicates that the peptide contains a T cell epitope.

However, only individual S.I. values above 2.0 (a response two-fold or greater over background) were used in calculating mean stimulation indices for each peptide for the group of patients tested. The results shown in Table II demonstrate that this patient (#466) responds very well to peptides RAE 7.D, RAE 69.1, RAE 64.1, and RAE 29.1. This indicates that these peptides contain *Amb a I.1* T cell epitopes recognized by T cells from this particular allergic patient.

TABLE II

10	Antigen	Concentration (mg/ml)	CPM	S.I.
	Medium	--	1677	--
	Amb a I.1	2	12113	
		10	25047	
15		50	45710	27.3
	RAE 22.E	5	3198	
		50	3581	2.1
20	RAE 7.D	5	3948	
		50	8572	5.1
	RAE 45.15	5	1160	
		50	1370	0.8
25	RAE 69.1	5	6800	
		50	11464	6.8
	RAE 70.1	5	4116	2.5
30		50	3888	
	RAE 65.1	5	1236	
		50	1784	1.1
35	RAE 64.1	5	12828	7.7
		50	2259	
	RAE 29.1	5	6998	
		50	13078	7.8

The above procedure was followed with a number of other patients. Individual patient results were used in calculating the mean S.I. for each peptide if the patient responded to the *Amb a* I.1 protein at an S.I. of 2.0 or greater and the patient responded to at least one peptide derived from *Amb a* I.1 at an S.I. of 2.0 or greater. A summary of positive experiments from 39 patients is shown in Fig. 8. The bar represents the cumulative rank of the peptide response in the 39 patients. To determine the cumulative rank, the 5 peptides with the highest S.I. in each patient were determined and assigned a numerical rank in descending order with 5 representing the strongest response. The ranks for each peptide were then summed in the 39 patients to determine the cumulative rank for the peptide. The number above each bar is the mean S.I. of the positive responses (S.I. of 2.0 or greater) from the group of patients to that peptide. In parentheses above each bar is the positivity index (P.I.). The P.I. for each peptide is determined by multiplying the mean S.I. by the percent of patients who responded to that peptide. The P.I. therefore represents both the strength of the response (S.I.) and the frequency of a response to a peptide in the group of patients tested. For example, peptide RAE 69.1 had the highest cumulative rank so it was the best peptide response in the overall population of 39 even though it did not have the highest mean S.I. Similarly, RAE 70.1 had the highest mean S.I. but not the best cumulative rank or P.I. Thus, the response to RAE 70.1 was strong when it occurred but it did not occur as frequently in the population as the response to other peptides. The peptide with the highest P.I., RAE 29.1, also had a strong S.I. and the second highest cumulative rank. The response to this peptide was therefore generally strong and relatively frequent in this population.

#### Example VII T Cell Epitope Fine Map Studies with *Amb a* I.1

Based on the above analysis 4 major areas of T cell reactivity within *Amb a* I.1 were identified. (Regions 1, 2, 3 and 4). All 39 patients responded to *Amb a* I.1 and a peptide from at least one of these regions: Region 1, amino acids residues 48-107; Region 2, amino acid residues 171-216; Region 3, amino acid residues 278-322; and Region 4, amino acid residues 331-377. Based in part on the T cell reactivity shown in Fig. 8, *Amb a* I.1 peptides were selected and modified by addition or deletion of amino acid residues at either the 5' or 3' end of the peptide. This set of peptides is shown in Fig. 9. T cell studies similar to those of Example VI were performed using these selected peptides to more precisely define the major areas of T cell reactivity within Regions 1-4 of the *Amb a* I.1 protein. For example, PBMC from a single ragweed-allergic patient were isolated as described in Example VI and were stimulated with 20 mg/ml of recombinant *Amb a* I.1 as described above. The results of proliferation assays with this one patient to selected peptides using irradiated (24,000 RADS) autologous EBV B cells as antigen presenting cells is shown in Table III. The data indicates that T cells from this patient respond well to RAE 7.D, RAE 70.1, RAE 40.1-4, and RAE 28.1-2.

- 36 -

TABLE III

	<u>Antigen</u>	<u>Concentration (mg/ml)</u>	<u>CPM</u>	<u>S.I.</u>
5	Medium	--	2762	--
	PHA	1	97864	35.4
	Amb a I.1	2	28025	
10		10	56172	
		50	86598	31.4
	RAE 22.E	5	3318	1.2
		50	784	
15	RAE 7.D	5	53292	
		50	60943	22.1
	RAE 40.1-4	5	4504	
20		50	13549	4.9
	RAE 69.1	5	5213	
		50	5981	2.2
25	RAE 70.1	5	14223	5.1
		50	4729	
	RAE 64.1	5	3418	1.2
		50	2538	
30	RAE 28.1-2	5	7542	
		50	24208	8.8

The above procedure was followed with a number of other patients and yielded 47 positive experiments. A summary of the results for Region 1 peptides of the *Amb a* I.1 protein is shown in Fig. 10, Region 2 peptides in Fig. 11, Region 3 peptides in Fig. 12, and Region 4 peptides in Fig. 13. As described in Example VI, the bar shows the cumulative rank. In these experiments the 3 best peptide responses were ranked. The number above each bar represents the mean S.I. for the peptide and the number in parentheses the P.I. for the peptide. In these experiments an individual S.I. of 2.5 or greater was used in calculating the mean S.I.

Fig. 10 indicates that the major area of T cell reactivity within Region 1 of the *Amb a* I.1 protein is represented by peptides RAE 6.D, RAE 7.D, RAE 40.1, RAE 40.1-4, and RAE 40.1-5. Thus, a preferred area of major T cell reactivity within Region 1 comprises amino acid residues 57-101. Fig. 11 indicates that there is a broad area of weak T cell reactivity in Region 2 of the *Amb a* I.1 protein relative to Region 1. A preferred area of major T cell reactivity within Region 2 thus comprises amino acid residues 182-216. Fig. 12 shows that the most frequent and dominant response within Region 3 of the *Amb a* I.3 protein is to peptide RAE 64.1. However, another area of T cell reactivity is represented by peptide RAE 66.1. A preferred area of major T cell reactivity within Region 3 comprises amino acid residues 280-322. Fig. 13 indicates that the major area of T cell reactivity in Region 4 is represented by peptides RAE 28.1-2, RAE 28.1-1, and RAE 28.1. From this analysis, a preferred area of major T cell reactivity within Region 4 comprises amino acid residues 342-377.

To further validate and define peptides derived from the *Amb a* I.1 protein comprising T cell epitopes, selected peptides from Regions 1-4 were further modified by addition or deletion of amino acid residues as described above. These selected modified peptides are shown in Fig. 14. To determine the T cell reactivity of these peptides, PBMC were stimulated with recombinant *Amb a* I.1 and assayed with the selected peptides as described in Example VI, except in some cases autologous PBMC (irradiated 3.500 RADS) were used as antigen presenting cells. The assay was followed with a number of patients and resulted in 23 positive experiments. In these assays an individual S.I. of 2.0 was used in calculating the mean stimulation index. In Figs. 15-18, the bar represents the cumulative rank of each peptide. The best 3 peptide responses for each patient were ranked as described above. The mean S.I. for each peptide and percent positive are found above the bar.

#### **Example VIII *Amb a* I.1 Specific T Cell Cross-Reactivity To Other *Amb a* I Family Members and *Amb a* II**

To determine the relationship of the *Amb a* I family members (i.e., *Amb a* I.1, I.2, I.3, and I.4) at the level of T cell reactivity, PBMC were stimulated with recombinant *Amb a* I.1 (r *Amb a* I.1) as described in Example VI. For assay,  $2 \times 10^4$  rested cells were restimulated in the presence of an equal number of autologous EBV-transformed B cells (irradiated with

25,000 RADS) with various concentrations (0-100 mg/ml) of recombinant *Amb a* I.1, I.2, I.3, I.4, *Amb a* II, or ragweed pollen extract in a volume of 200 ml complete medium in duplicate or triplicate wells in 96-well round bottom plates for 3 days. Each well then received 1 mCi tritiated thymidine for 16-20 hours. The counts incorporated were assessed and analyzed as described in Example VI. Table IV shows the S.I., P.I., percent of cultures positive for the assay antigen, and number of cultures analyzed. For these assays an individual S.I. greater or equal to 2.0 was used in calculating the mean S.I. *Amb a* I.1 stimulated cells respond less well and less frequently to the least homologous family members I.3 and I.2. However, this level of cross-reactivity, 68.8% and 60% respectively, is still considered high. The lowest level of *Amb a* I.1 stimulated T cell cross-reactivity is noted to the least homologous allergen, *Amb a* II. In addition, *Amb a* I.1 reactive T cells are found at a high percentage from ragweed pollen extract stimulated cultures, again demonstrating the importance of the *Amb a* I.1 allergen.

TABLE IV

Assay Antigen	Recombinant <i>Amb a</i> I.1 Specific Lines			
	S.I.	% positive	n	P.I.
Pollen Extract	13.5	87.8%	41	1185
r <i>Amb a</i> I.1	27.5	95.1%	41	2616
r <i>Amb a</i> I.2	5.0	60%	5	300
r <i>Amb a</i> I.3	9.3	68.8%	16	639
r <i>Amb a</i> I.4	8.5	100%	11	850
r <i>Amb a</i> II	12.8	55.2%	29	70



**Example IX T Cell Cross-Reactivity of *Amb a* I.1 or *Amb a* I.3 Stimulated Cells to *Amb a* I.1 or *Amb a* I.3 Peptides**

Since a high degree of T cell cross-reactivity could be demonstrated between *Amb a* I.1 and the most highly sequence divergent family member *Amb a* I.3, the ability of *Amb a* I.1 stimulated cells to recognize both *Amb a* I.1 peptides and homologous *Amb a* I.3 peptides was assessed. In a similar fashion, the ability of *Amb a* I.3 stimulated cells to recognize both *Amb a* I.3 peptides and homologous *Amb a* I.1 peptides was determined. The peptides used in this study are listed below. The sequences for the *Amb a* I.1 homologous *Amb a* I.3 peptides were based on an alignment of the *Amb a* I.3 sequence to the *Amb a* I.1 sequence.

***Amb a* I.1 peptides**

AMB 1-2.1 KGTVGGKDGDIYTVTSELDDDDVAN  
 AMB 1-4.1 AENRKALADCAQGFSGKGTVGGKDG  
 AMB 2-1.1 GPAAPRAGSDGDAISISGSSQ  
 AMB 3-1.1 GSYAIGGSASPTILSQGNRFCAPDERSK  
 AMB 3-4.1 FFQVVNNNYDKWGSYAIGGSASPT  
 AMB 4-3.1 VLENGAIFVASGVDPVLTPEQSAGMIP

***Amb a* I.3 peptides**

AMB 1-2.15 KGTYGGKWGDVYTVTSNLDDDDVAN  
 AMB 1-4.15 ENNRQALADCAQGFAGKGTYGGKWGD  
 AMB 2-1.15 GPILRQASDGDITINVAGSSQ  
 AMB 3-1.15 GTYAIGGSSAPTILCQGNRFLAPDDQIK  
 AMB 3-4.15 FFQVVNNNYDRWGTYAIGGSSAPT  
 AMB 4-3.15 LLENGAIFVTSGSDPVLTPVQSAGMIP

PBMC from ragweed allergic patients were stimulated with 20 mg/ml recombinant *Amb a* I.1 protein as described in Example VI with the addition of cultures of PBMC which were stimulated with 20 mg/ml recombinant *Amb a* I.3 protein. Assays were performed as described in Example VI except the homologous *Amb a* I.3 peptides were also tested at various doses. In these experiments an individual S.I. of 2.0 or greater was used in calculating the mean S.I. Fig. 19 shows the results from a set of 9 matched patients stimulated with either recombinant *Amb a* I.1 protein or recombinant *Amb a* I.3 protein and tested on the set of *Amb a* I.1 peptides described above. Each bar represents the P.I. The dark bar represents cells stimulated with recombinant *Amb a* I.1 protein and tested with *Amb a* I.1 peptides, whereas, the stippled bar represents cells stimulated with recombinant *Amb a* I.3 protein and tested with *Amb a* I.1 peptides. With two exceptions, the results directly

parallel each other, indicating that cells stimulated with recombinant *Amb a* I.3 proteins recognize *Amb a* I.1 derived peptides comprising at least one T cell epitope. One exception is that cells primed with *Amb a* I.3 protein recognize the RAE 7.D peptide poorly compared to recombinant *Amb a* I.1 stimulated cells. In addition, cells stimulated with recombinant *Amb a* I.3 protein gave stronger responses overall to *Amb a* I.1 derived peptides from Region 4 than the recombinant *Amb a* I.1 stimulated cells, particularly peptide RAE 28.1-2.

Cells were also stimulated with either recombinant *Amb a* I.1 or recombinant *Amb a* I.3 and tested for reactivity with *Amb a* I.3 derived peptides (Fig. 20). The results indicate that cells stimulated with recombinant *Amb a* I.1 protein recognize the *Amb a* I.3 derived peptides in a similar pattern to that of cells stimulated with recombinant *Amb a* I.3 protein. In Fig. 20 the bars represent the P.I. and the number above the bar the mean S.I. in the nine patients. The ".15" designation following the peptide name indicates that the peptide is derived from the *Amb a* I.3 protein. The data suggests a high degree of cross-reactivity at the T cell level between the *Amb a* I.1 protein and the *Amb a* I.3 protein.

#### **Example X IgE Analysis With Peptides Derived From *Amb a* I.1**

To analyze IgE reactivity of peptides derived from the *Amb a* I.1 protein, a direct binding ELISA was performed according to the procedure described in Example III. The source of IgE for this analysis was a pool of ragweed allergic patient plasma from 38 ragweed skin test positive patients. The ELISA protocol was the same as Example III except that the antigen coating with peptides and proteins was a concentration of 10 mg/ml at 100 mls/well. Fig. 21 shows the graphed results of this assay demonstrating strong reactivity to both SPE and biochemically purified *Amb a* I.1. By this assay there is no detectable binding to any of the peptides. A set of these assays was also run with rabbit and mouse antisera to demonstrate that the coating of the peptides onto these plates was successful.

#### **Example XI Further T Cell Epitope Fine Map Studies With *Amb a* I.1**

To further define the regions of *Amb a* I.1 specific T cell reactivity, selected peptides from Figure 14 were analyzed. *Amb a* I.1 specific T cells lines were derived from 28 ragweed allergic patients as described in Example VI. These lines were assessed for their ability to proliferate in response to the peptide in the presence of autologous EBV transformed antigen presenting cells by the uptake of tritiated thymidine as described previously. Several lysine substituted peptides derived from the peptide RAE 70.1 sequence were designed to increase the solubility of the peptide from Region 2. These peptides shown in Figure 22 were tested against non-substituted controls to determine if the modification resulted in a change in T cell reactivity. In addition, truncations of peptide AMB 4-6.1 (amino acid residues 347-377) were tested to further refine the relative T cell reactivity in that region.

Figure 22 shows the responses of the 28 T cell lines to these peptides as analyzed by the positivity index, the mean stimulation index and the percent of positive responses. The positivity index, as defined previously, is the mean stimulation index multiplied by the percent of patients responding to a peptide. Responses to peptides were considered positive if they were greater or equal to 2 fold over background. Figure 22 demonstrates that certain lysine substituted peptides in the 182-216 sequence resulted in greater T cell responses, indicating that not all substitutions are recognized equally by a given T cell line. In addition, the T cell responses to the lysine modified peptides having a substitution of cysteine at position 212 with serine reflect the responses of T cell lines tested with peptides without the lysine substitutions. Thus, the decrease in response to AMB 2-10.1, with a substitution of leucine for cysteine at 212, reflects the decrease in the percentage of patients responding to this peptide relative to peptides AMB 2-9.1 and AMB 2-11.1 which contain serine and glutamic acid substitutions, respectively, at position 212.

In addition, the truncation peptides derived from peptide AMB 4-6.1 showed indistinguishable responses based on this analysis. Thus, the shortest sequence, 345-370 was chosen for further analysis. Substitutions at positions 360 and 361 were made to increase the stability of the AMB 4-9.1 peptide which contained an acid-sensitive aspartic acid-proline bond. As shown in Figure 23, three peptides were synthesized in which the proline at position 361 was substituted with either alanine, serine or glycine (AMB 4-9.1DA (SEQ ID NO:147), AMB 4-9.1DS (SEQ ID NO:148), and AMB 4-9.1DG (SEQ ID NO:149), respectively). Similarly, five peptides were synthesized in which the aspartic acid at position 360 was substituted with the following: glutamic acid (AMB 4-9.1EP, SEQ ID NO:142); asparagine (AMB 4-9.1NP, SEQ ID NO:143); alanine (AMB 4-9.1AP, SEQ ID NO:144); serine (AMB 4-9.1SP, SEQ ID NO:145); and glutamine (AMB 4-9.1QP, SEQ ID NO:146). This group of modified peptides were tested to determine whether a change in T cell reactivity occurred. 28 *Amb a* I.1 specific T cells lines were analyzed by the relative strength of the response to the modified peptides in a given T cell line compared to the overall response to all peptides from that line. Figure 23 shows the ranked sum of the strongest three peptide responses in the 28 T cell lines. The ranked sum was determined as described previously with the strongest response to a peptide given a value of 3, the second strongest a 2, and the third strongest a value of 1. The values from all 28 lines were then added to obtain the ranked sum. Figure 23 indicates that the modified peptide AMB 4-9.1DA elicits T cell responses similar to those of the native sequence AMB 4-9.1. In contrast, other substitutions, while eliciting strong responses do not rank as highly in this analysis.

Based in part on the above described analyses, five peptides were selected for further study (AMB 1-2.1, AMB 2-6.1, AMB 2-9.1, AMB 3-5.1 and AMB 4-9.1). An additional peptide, RA-02.1 (SEQ ID NO:150) was also synthesized and tested to confirm that the serine substitution at position 212 was recognized by *Amb a* I.1 specific T cell lines. The T cell reactivity of these peptides in 32 different *Amb a* I.1 specific T cell lines was determined

as described previously. Figure 24 demonstrates the relative positivity indices, percent of positive responses and mean stimulation indices of these six peptides compared to the overall response to *Amb a* 1.1. This figure shows that peptide AMB 4-9.1 elicits the strongest response in a high percentage of patients.

5 In contrast to previous data, the response to the 182-216 peptide (AMB 2-6.1) and substituted variants within this sequence (RA-02.1) gave weaker stimulation indices. This could reflect a potential toxic effect of the peptides in the T cell assays. To investigate this possibility, an assay was conducted to determine the ability of the peptides to inhibit T cell proliferation. Inhibition was defined as proliferation of a T cell line in response to the  
10 peptide which was less than half the proliferation of media control T cell lines plus autologous EBV transformed antigen presenting cells. As seen in Table V, the percent of patients in which inhibition of T cell proliferation is seen is far greater in cultures where the AMB 2-6.1, RA-02.1, and AMB 2-9.1 peptides are present as compared to AMB 1-2.1 or  
15 AMB 4-9.1. This data suggests that AMB 2-6.1, RA-02.1 and AMB 2-9.1 may be inhibitory due to toxicity in *in vitro* T cell line culture. This is further supported by the finding that these peptides also inhibit a T cell line from patient 956.2 which is specific for an irrelevant antigen, native *Fel d* I, as shown in Figure 26.

TABLE V

20

ASSESSMENT OF PEPTIDE TOXICITY IN *Amb a* 1.1  
PRIMED SECONDARY T CELL CULTURE

25	peptide	#/(%)patients	low dose only	high dose only	both
	AMB 1-2.1	3/46 (7)	2/3	1/3	0
	AMB 2-6.1	14/44 (32)	0	11/14	3/14
30	RA-02.1	19/45 (41)	0	15/19	4/19
	AMB 2-9.1	23/38 (61)	0	14/23	9/23
	AMB 4-9.1	1/46 (2)	0	0	1/1
35	AMB 3-5.1	19/46(41)	0	17/19	2/19

Legend: Toxicity defined as proliferation, as assessed by incorporation of tritiated thymidine, which is less than half the proliferation of media control of an *Amb a* 1.1

stimulated T cell line plus APC. Low dose = 5 mg/ml peptide in secondary assay, high dose = 50 mg/ml peptide.

5 A peptide from Region 2, RAE 70.1 (SEQ ID NO:44) was modified to minimize dimerization via disulfide linkages by substituting serine for cysteine at position 212. In addition, peptide RAE 70.1 was divided into two fragments and modified to increase solubility and to further define the residues necessary for T cell reactivity. As shown in Figure 25, a fragment of RAE 70.1 (amino acid residues 194-216) was modified by substitution or addition of amino acids which would increase the hydrophilicity and decrease the pI of the peptide to thereby increase solubility.

10 Peptides Amb 2-18.1 (SEQ ID NO:126), Amb2-19.1 (SEQ ID NO:127), Amb2-20.1 (SEQ ID NO:128) and Amb2-21.1 (SEQ ID NO:129) correspond to different lengths of the original peptide which were synthesized (including the serine substitution at position 212) and analyzed to determine solubility. It was found that peptide Amb2-19.1 (residues 200-217) was most soluble. Thus, substitutions and/or amino acid additions to this peptide were made in order to further increase the solubility while maintaining T cell reactivity. For example, in peptides Amb2-22.1 (SEQ ID NO:130) and Amb2-23.1 (SEQ ID NO:131) isoleucine at position 201 was substituted with glutamic acid (Amb2-22.1) or lysine (Amb2-23.1) to lower the pI of the resulting peptide and avoid precipitation of the peptide at physiological pH. Similarly, the following peptides were also synthesized with various substitutions or additions designed to decrease the pI and increase the hydrophilicity of the peptide, each is shown in Figure 25: Amb2-26.1 (SEQ ID NO:132); Amb2-28.1 (SEQ ID NO:133); Amb2-30.1 (SEQ ID NO:134); Amb2-32.1 (SEQ ID NO:135); Amb2-33.1 (SEQ ID NO:136); Amb2-34.1 (SEQ ID NO:137); Amb2-35.1 (SEQ ID NO:138); Amb2-36.1 (SEQ ID NO:139); Amb2-37.1 (SEQ ID NO:140); and Amb2-38.1 (SEQ ID NO:141).

25 Figures 27 and 28 show representative examples of proliferation of two individual *Amb a* I.1 specific T cell lines to peptides selected from those shown in Figure 25. Assays were performed as described previously. Both Figure 27 and Figure 28 indicate a hierarchy of responses to these peptides. Patient 119 shown in Figure 27, has a hierarchy of response from strongest to weakest as follows: AMB2-23.1>AMB2-22.1>AMB2-30.1>AMB2-26.1>AMB2-33.1>AMB2-32.1>AMB2-18.1>AMB2-19.1. Peptides AMB2-34.1, AMB2-35.1, and AMB 2-9.1 did not elicit significant T cell proliferation in this patient. Patient 1199 shown in Figure 28 demonstrated strong responses to AMB 2-26.1>AMB 2-22.1>AMB 2-33.1. T cells from this patient showed less than 2 fold over background responses to all other peptides. These responses are not significantly different from the media control of the T cell line plus autologous EBV antigen presenting cells.

35 To more closely examine the T cell epitopes within the above set of peptides to which *Amb a* I.1 specific T cell lines respond, a T cell clone was generated by limiting dilution from an *Amb a* I.1 specific T cell line stimulated with AMB 2-10.1. Briefly, an *Amb a* I.1 specific

T cell line from patient 776 was shown to respond to AMB 2-10.1 in a proliferation assay as described previously. To generate a AMB 2-10.1 specific T cell clone, *Amb a* I.1 specific T cells were plated at 0.3 cells/well in a V-bottom 96 well plate with 20,000 irradiated autologous EBV transformed antigen presenting cells, AMB 2-10.1 at 40 mg/ml, leukoagglutinin at 1 mg/ml, and recombinant human IL-2 and IL-4 at 10 units/ml in a total volume of 100ml/well. Plating the T cells at 0.3 cells/well insures not more than 1 T cell/well can potentially proliferate to the peptide with the progeny of that T cell representing a clonal population. After five days, wells received an additional 15 units/ml of recombinant human IL-2 and IL-4 to expand peptide-specific T cells. This addition of IL-2 and IL-4 was repeated again every three days for the duration of the culture period. Twelve days after the initiation of culture, T cells in the wells were restimulated with 20,000 irradiated EBV antigen presenting cells and 40 mg/ml AMB 2-10.1. This was repeated again 20 days after the initial stimulation. T cells from wells which showed signs of growth were separated from cell debris by density centrifugation as described previously in the generation of T cell lines. The T cell clone was then expanded with additional IL-2 and IL-4 at 10 units/ml until there were significant numbers to assay for proliferation.

Assessment for T cell proliferation to selected peptides from those shown in Figure 25 was performed as described previously for T cell lines. The results of a representative assay using this AMB 2-10.1 specific T cell clone are shown in Figure 29. The data indicates that the T cell epitope recognized by this clone is contained by all the peptides except AMB 2-34.1 and AMB 2-35.1. Thus, a substitution of the tryptophan at position 208 eliminates the ability of the T cell clone to respond to these peptides. This data suggests that substitutions on the N and C terminal ends of the truncated peptides does not effect the recognition of the peptides by the T cell clone and, thus, does not alter the residues which comprise the T cell epitope.

#### **Example XII Histamine Release Analysis With Purified *Amb a* I.1 and With Peptides Derived From *Amb a* I.1**

The objective of the histamine release analysis was to compare the effects of *Amb a* I.1 or *Amb a* I.1-derived peptides in an *in vitro* allergic response system. The release of histamine via IgE recognition and IgE receptor crosslinking on viable cells directly assays the allergic potential of a protein antigen.

The histamine release assay used for these studies is based on the detection of an acylated derivative of histamine using a specific monoclonal antibody (Morel, A.M. and Delaage, M.A. (1988) *J. Allergy Clin. Immunol.* 82:646-654). The assay was performed in two steps: 1) the release of histamine from basophils present in heparinized whole blood in the presence of different concentrations of protein or peptide; and 2) the assay of histamine present in the supernatants of the release reactions following cell removal by centrifugation.

The reagents for this second step are available commercially as a competitive radioimmunoassay from AMAC Inc. (Westbrook, ME).

Heparinized whole blood was drawn from ragweed allergic patients. The test protein Amb a I.1 and peptides AMB 1-2.1, AMB 2-9.1, AMB 4-9.1, and AMB 3-5.1 were each  
5 diluted to 2X the final release concentration in PACM buffer (PIPES 25mM, NaCl 110mM, KCl 5.0 mM, human serum albumin 0.003% (w/v), CaCl<sub>2</sub> 5mM, MgCl<sub>2</sub> 2mM, pH7.3) and 0.2 ml of each dilution was added to a 1.5 ml polypropylene tube. A 0.2ml aliquot of blood was then added to each tube and the release reactions started by inversion of the tubes. A negative control of 0.2 ml blood incubated with 0.2 ml PACM buffer was also included. The  
10 release reactions were performed for 30 minutes at 37°C. The tubes were then centrifuged in a microfuge at 1500 RPM for 3 minutes and the supernatants were carefully removed and analyzed or stored at -20°C for later analysis. To analyze the total histamine released, 0.1 ml blood was diluted with 0.9 ml PACM buffer and boiled 3 minutes. This tube was then centrifuged 2 minutes at 12,000 RPM and the supernatant was removed and save for analysis.

15 For the competition radioimmunoassay, a 50 ml aliquot of each release supernatant was mixed with 150 ml of histamine release buffer (supplied with the kit from AMAC). The diluted supernatant (100 ml) was added to a kit tube containing an acylation reagent. A 50 ml aliquot of acylation buffer was then immediately added and the tube mixed by vortexing. The acylation reactions were incubated at least 30 minutes at room temperature. A set of  
20 histamine standards supplied with the kit were acylated at the same time. 50 ml of each acylation reaction was then placed in a tube coated with a monoclonal antibody which specifically recognizes the acylated form of histamine. A 0.5 ml aliquot of <sup>125</sup>I-labelled tracer was added and the tubes were incubated at least 18 hours at 4°C. The solutions were aspirated from the tubes and the radioactivity bound to the tube was measured on a gamma  
25 counter (Cobra 5005, Beckman, Inc.) for two minutes per tube.

A standard curve was generated from the histamine standard counts and graphed on a semi-log plot. Since this is a competitive assay (the <sup>125</sup>I-labelled tracer competes with acylated histamine in the samples for binding to the antibody-coated tubes), the lower the number of radioactive counts measured, the greater the amount of histamine in the sample.  
30 The amount of histamine in each sample data point was extrapolated from the standard curve using a computer statistical program (StatView for the MacIntosh). The assay is sensitive to 0.2 nM histamine.

To study whether peptides derived from the Amb a I.1 sequence could induce histamine release, blood samples from 8 ragweed-allergic patients who exhibited histamine  
35 release to native Amb a I.1 were analyzed with AMB 1-2.1, AMB 2-9.1, AMB 4-9.1 and AMB 3-5.1. Six or more 10-fold dilutions of Amb a I.1 starting at 10 mg/ml (approximately 0.66 mM) were analyzed. Five five-fold dilutions of each peptide were also analyzed (ranging from 50 mg/ml to 0.08 mg/ml, this is approximately 15 mM to 24 nM in concentration for each peptide). The peptide concentrations were selected to encompass the

higher end of the concentration curve. Figure 30 shows the representative results from 1 of the patients (#1273). The graph shows the concentration of each antigen in mg/ml versus the percent of total histamine released. The results are presented as the percent of the total histamine released for that patient, since patients vary greatly in their overall releasable histamine levels. This variability is likely to result from variation in the number of basophils per ml between patients and the histamine content of each basophil. As Figure 30 shows, *Amb a* I.1, at all the concentrations tested, yields a high level of histamine release in this patient. Similar experiments using blood samples from other patients have shown that for some patients the *Amb a* I.1 concentration does not become limiting until  $10^{-5}$ - $10^{-6}$  mg/ml. By contrast, there is no discernible histamine release to any of the four *Amb a* I.1-derived peptides which were used. Similar results were obtained for seven other patients similarly tested.

In another experiment, plasma samples from 94 ragweed-allergic individuals, who tested positive for IgE binding to *Amb a* I.1 were analyzed for IgE binding to peptides Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1 by direct binding ELISA as described above. Plasma samples were studied at a 1:3 and 1:30 dilution. None of the 94 patients tested had measurable IgE binding to either Amb 1-2.1 or Amb 2-36.1 (data not shown). Six patients had a low level of IgE binding to Amb 4-9.1 (data not shown). These results indicate that approximately 7% of ragweed-allergic patients have low plasma IgE levels to Amb 4-9.1

In order to assess the relative contribution of Amb 4-9.1-specific IgE to the overall level of IgE to *Amb a* I.1, plasma from one patient with IgE to Amb 4-9.1 was depleted of IgE to Amb 4-9.1 by repeatedly absorbing the plasma on plates coated with Amb 4-9.1-03 conjugated ovalbumin. A mock absorption was performed in parallel using plates coated with Amb 1-2.1 conjugated to ovalbumin. Previous experiments have shown that a somewhat higher O.D. signal can be obtained when the ovalbumin conjugate is used for plate coating. After absorption, the original, non-absorbed plasma samples were analyzed for IgE binding to *Amb a* I.1 in a direct assay. The efficiency of absorption was verified by also analyzing binding to Amb 4-9.1. The results indicated that absorption with Amb 4-9.1 efficiently removed any detectable binding to Amb 4-9.1. The mock absorption with 1-2.1 yielded a slight decrease in the O.D. signals seen in the unabsorbed sample, as indicated by the shift curve for Amb 1-2.1 absorbed plasma to the left of the curve for unabsorbed plasma. The curve for Amb 4-9.1 shifted very slightly to the left of the Amb 1-2.1-absorbed plasma curve. This result suggests that IgE binding to Amb 4-9.1 represents a minor but measurable component of the overall level of IgE to *Amb a* I.1.

To further assess the relative contribution of anti-AMB 4-9.1-03 IgE to the overall anti-*Amb a* I.1 response, a competition ELISA was performed using Amb 4-9.1 and *Amb a* I.1 in solution to compete for binding to *Amb a* I.1. For a negative control, competition was also performed using Amb 1-2.1. This experiment shows that Amb 4-9.1 does not detectably compete for IgE binding to *Amb a* I.1 at concentrations where very effective competition is



seen with soluble *Amb a* I.1. The results (data not shown) indicate that the change in IgE binding detected to Amb 4-9.1 when the plasma IgE response is competed with *Amb a* I.1 and Amb 4-9.1. As expected, incubation of the plasma sample with soluble Amb 4-9.1 can compete for binding to Amb 4-9.1. However, soluble *Amb a* I.1 competes much more efficiently since much lower concentrations (in  $\mu\text{g/mL}$ ) of *Amb a* I.1 protein than Amb 4-9.1 peptide are required to compete for binding to Amb 4-9.1. This difference is even more distinct if the comparison is made on an equimolar basis since the peptide is in greater than ten-fold excess relative to *Amb a* I.1 at equivalent  $\mu\text{g/mL}$  concentrations. This result suggests that in this patient sample, the portion of the IgE response detected which is directed to an epitope defined by the Amb 4-9.1 peptide sequence has lower affinity for Amb 4-9.1 than for *Amb a* I.1.

All peptide candidates were also examined for their ability to release histamine *in vitro* from basophils of ragweed-allergic patients. The objective of the histamine release analysis was to measure the effects of *Amb a* I.1 peptides Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1 *in vitro* allergic response system. Heparinized whole blood from ragweed-allergic patients was incubated with different concentrations of each peptide (at 0.08-50  $\mu\text{g/mL}$ ) or *Amb a* I.1 (at  $10^{-7}$  to 10  $\mu\text{g/mL}$ ). The levels of histamine release were measured using commercially available radioimmunoassay (Amac, Inc., Westbrook, ME). No patient with histamine release to native *Amb a* I.1 ( $n=19$ ), released histamine to peptides Amb 1-2.1 ( $n=19$ ), Amb 2-36.1 ( $n=19$ ) and Amb 4-9.1 ( $n=14$ ) (data not shown). Histamine release to Amb 4-9.1 was also not detected when blood from two patients with anti Amb 4-9.1 antibodies was analyzed (data not shown). The failure of Amb 4-9.1 to induce histamine release in two patients with IgE to Amb 4-9.1 is possibly due to the inability of the peptide to mediate the crosslinking of the IgE receptors on blood basophils.

#### **Example XIIT Cell Responses to Ragweed Peptides**

Peripheral blood mononuclear cells (PBMC) were purified by lymphocyte separation medium (LSM) centrifugation of 60 ml of heparinized blood from ragweed-allergic patients who exhibited clinical symptoms of seasonal rhinitis and were skin prick test positive for ragweed. T cell lines were established from these cells by stimulation of  $1-2 \times 10^6$  PBMC/ml in RPMI-1640 containing 5% human AB serum (complete medium) with native *Amb a* I.1 at 20 mg/ml for 5-6 days at 37°C in a humidified CO<sub>2</sub> incubator. Viable cells were purified by LSM centrifugation and cultured in complete medium supplemented with 5 units recombinant human IL-2/ml and 5 units recombinant human IL-4/ml for up to three weeks until the cells no longer responded to lymphokines and were considered "rested". The ability of the T cells to proliferate to *Amb a* I.1 sequence-derived synthetic peptides was then assessed.

For assay,  $2 \times 10^4$  rested cells were restimulated in the presence of  $2 \times 10^4$  autologous Epstein-Barr virus (EBV)-transformed B cells (gamma-irradiated with 25,000

RADS) or  $5 \times 10^4$  autologous PBMC (3,500 RADS) with various concentrations of *Amb a* I.1 synthetic peptides in a volume of 200 ml complete medium in duplicate or triplicate wells in 96-well round bottom plates for 3 days. Each well then received 1 mCi tritiated thymidine for 16-20 hours. The counts incorporated were collected onto glass fiber filter mats and  
5 processed for liquid scintillation counting. Table III shows the results of a representative assay. The maximum response in a titration of each peptide is expressed as the S.I. or stimulation index. The S.I. is the CPM incorporated by cells in response to peptide divided by the CPM incorporated by cells in medium only. An S.I. value greater than the background level is considered "positive" and indicates that the peptide contains a T cell epitope.  
10 However, only individual S.I. values above 2.0 (a response two-fold or greater over background) were used in calculating mean stimulation indices for each peptide for the group of patients tested. The results shown in Table VI demonstrate that this patient (#1715) responds very well to peptides RAE 74.1, Amb 4-9.1, RAE 61.1, RAE 62.1. This indicates that these peptides contain *Amb a* I.1 T cell epitopes recognized by T cells from this  
15 particular allergic patient.

TABLE VI

	<u>Antigen</u>	<u>Concentration (mg/ml)</u>	<u>CPM</u>	<u>S.I.</u>
5	Medium	--	452	1
	Amb a I.1	10	47822	
		100	63494	139.9
10	RAE 61.1	5	767	
		50	5910	13.0
	RAE 80.1	5	640	
		50	1067	2.4
15	RAE 45.1	5	589	
		50	1106	2.4
	RAE 75.1	5	1094	2.4
20		50	521	
	RAE 62.1	5	2017	
		50	4923	10.8
25	RAE 65.1	5	1048	2.3
		50	462	
	RW 31	5	892	
		50	1309	2.9
30	RAE 74.1	5	18992	41.8
		50	10633	7.8
	AMB 4-9.1	5	14065	31.0
35		50	8018	

The above procedure was followed with a number of other patients. Individual patient results were used in calculating the mean S.I. for each peptide if the patient responded to the *Amb a* 1.1 protein at an S.I. of 2.0 or greater and the patient responded to at least one peptide derived from *Amb a* 1.1 at an S.I. of 2.0 or greater. A summary of positive experiments from 57 patients is shown in Fig. 31. The bar represents the cumulative rank of the peptide response in the 57 patients. To determine the cumulative rank, the 5 peptides with the highest S.I. in each patient were determined and assigned a numerical rank in descending order with 5 representing the strongest response. The ranks for each peptide were then summed in the 57 patients to determine the cumulative rank for the peptide. The number above each bar in parenthesis is the percentage of the positive responses (S.I. of 2.0 or greater) from the group of patients to that peptide. In parentheses above each bar is the mean S.I. of the positive responses. The cumulative rank sum represents both the strength of the response (S.I.) and the frequency of a response to a peptide in the group of patients tested. For example, peptide AMB 4-9.1 had the highest cumulative rank so it was the best peptide response in the overall population of 57 even though it did not have the highest mean S.I. Similarly, RAE 27.1 had the highest mean S.I. but not the best cumulative rank. Thus, the response to RAE 27.1 was strong when it occurred but it did not occur as frequently in the population as the response to other peptides. Based on the data as shown in Fig. 31, peptides AMB 4-9.1, RAE 74.1, AMB 3-1.1, AMB 2-36.1, RAE62.1 and AMB 1-2.1 were identified as the peptides with the strongest T cell reactivity

#### Example XIV Selection of Peptides For Use in Human Therapy

To determine whether a peptide (candidate peptide) or a combination of candidate peptides are likely to comprise a sufficient percentage of T cell epitopes of ragweed pollen protein to induce T cell non responsiveness in a substantial percentage of a population of individuals sensitive to ragweed pollen protein and therefore, be a suitable composition for human therapy, the following evaluation is used. Candidate peptides are selected based, at least in part on the mean human T cell stimulation index for each of the candidate peptides in the set of peptides tested and the positivity index (the mean S.I. multiplied by the percent of positive responses) of the candidate peptides in the set of peptides tested. For example, the results in Example XIII (Fig. 31) indicate a number of peptides which contain strong T cell reactivity, and three of such peptides, Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1, were selected as candidate peptides for use in human therapy. The human T cell stimulation index for each peptide tested including the stimulation index for the candidate peptides is summed. For each individual, the human T cell stimulation index for the candidate peptides are divided by the sum of the human T cell stimulation indices of the remaining peptides in the set of peptides tested to determine a percent of T cell reactivity as shown below:

% of T cell Reactivity of candidate peptides =

$$\frac{\text{Sum of S.I.s for Each Candidate Peptide}}{\text{Sum of S.I.s of the set of overlapping peptides}} \times 100$$

- 5 A more liberal estimate of reactivity elicited by the candidate peptide, accounting for the overlaps giving rise to the total can be calculated as follows:

% of T cell Reactivity of Candidate Peptides =

$$\frac{N_t \text{ flanking peptide S.I.} + \text{Candidate S.I.} + C_t \text{ flanking peptide S.I.}}{\text{Sum of S.I.s from overlapping peptides}} \times 100$$

10

In this formula, "N<sub>t</sub> flanking peptide" refers to a peptide which comprises amino acid residues which overlap with amino acid residues located at the N-terminus of the candidate peptide in the amino acid sequence of the protein antigen from which the peptide is derived; "C<sub>t</sub> flanking peptide" refers to a peptide which comprises amino acid residues which overlap with amino acid residues located at the C-terminus of the candidate peptide in the amino acid sequence of the protein antigen from which the peptide is derived. In this calculation, stimulation indices for the candidate peptide, the N-terminal flanking peptide and the C-terminal flanking peptide are added and divided by the sum total of the stimulation indices for the entire set of overlapping peptides to obtain a percent of T cell reactivity for the candidate peptides. If a combination of two or more candidate peptides is selected each of which contains amino acid residues which overlap, this calculation cannot be used to determine a percent of T cell reactivity for each candidate separately. However this is not the case for selected candidates, Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1.

15

20

25 The values obtained for the percentage of T cell reactivity for the candidate peptides in each individual tested can be expressed as a range of lower and higher values of the results of the above described calculations. The mean percentage of T cell reactivity elicited by the candidate peptides can then be determined by averaging the values obtained for the individual responses to the candidate peptides.

30 For candidate peptides Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1, secondary *Amb a 1.1* T cell cultures derived from 57 ragweed-allergic subjects were analyzed for reactivity to an overlapping set of Amb a 1.1 peptides, and the highest stimulation index greater than or equal to 2.0 in response to each peptide was recorded for each individual tested. The data were then analyzed by the equation above for each individual tested, and the mean percentage of T cell reactivity was determined by averaging all the values obtained for the individual responses to the candidate peptides, Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1, with the following results.

35

Candidate Peptides	Range of mean percentage of T cell reactivity	Frequency of response to at least one peptide
Amb 1-2.1 + Amb 2-36.1 + Amb 4-9.1	26-35%	91%

These values meet the criteria for a combination of candidate peptides described earlier.

## 5 Example XV Administration of Peptides to Humans for Treatment of Allergy to Ragweed

### A. Composition of Phase I Formulation

For Phase I clinical trials, the drug product was a multi-peptide formulation comprising three peptides, Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1, which together were determined to cover a sufficient percentage of T cell epitopes of a population of individuals as discussed in Example XIV. Each peptide was purified to homogeneity (at least 97% pure) by known methods and a multi-peptide formulation was prepared in accordance with procedures discussed earlier. The multi-peptide formulation used in this Phase I clinical Study was in the form of a freeze-dried powder cake comprising 1500 µg/L (when reconstituted) of each of the peptides, Amb 1-2.1, Amb 2-36.1 and Amb 4-9.1, in a single vial. The excipients used were 5% mannitol and 0.05 M sodium phosphate buffer system in a single vial. The formulation was reconstituted just prior to use with sterile water for injection resulting in a solution with a pH of ca. 7.5. Normal saline (0.9%) was used for any dilutions beyond initial reconstitution.

### 20 B. Human Phase I Clinical Studies

This study was a double blind placebo-controlled safety study of thirty-six patients conducted at Johns Hopkins Asthma and Allergy Center Baltimore. Twenty-four patients received active therapy while twelve received placebo.

After selection by demonstration skin test sensitivity to ragweed extract, the patients were tested for allergic sensitivity to study medication (active or placebo) prior to the first treatment by intracutaneous tests with increasing concentrations of the therapy. first Cohort (Cohort A) of sixteen patients received increasing weekly doses of 7.5, 75, 250, 750, and 1500 µg (per peptide) of the multi-peptide formulation and four received placebo for five weeks. Cohort B of ten patients (6 active, 4 placebo) initiated a five week course of weekly fixed dose therapy (250 µg per peptide or placebo). After all patients in Cohort A completed the fourth week of therapy,, a third Cohort (Cohort C) of ten patients (6 active, 4 placebo) began a five-week course of weekly fixed dose therapy (750 µg per peptide or placebo). All patients underwent skin testing to ragweed extract and study medication (active or placebo) three weeks post treatment. Physical evaluation and clinical laboratory safety studies were done at baseline and at eight weeks.

Preliminary data indicates that all doses were tolerated and the multi-peptide formulation appears to be safe. Although this study was designed to evaluate safety, preliminary data suggests that the multi-peptide formulation may possibly be reducing late phase responses to ragweed allergen in the skin. Further statistical analysis of the Phase I Clinical studies are pending. Phase II Clinical Studies are pending in the United States and Canada

Equivalents:

Although the invention has been described with reference to its preferred embodiments, other embodiments can achieve the same results. Variations and modifications to the present invention will be obvious to those skilled in the art and it is intended to cover in the appended claims all such modifications and equivalents that follow in the true spirit and scope of this invention.

- 54 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

5

(i) APPLICANT: Kuo, Mei-chang  
Garman, Richard  
Greenstein, Julia  
Evans, Sean  
10 Amsberry, Kent  
Shaked, Ze'ev

(ii) TITLE OF INVENTION: T CELL EPITOPES OF THE MAJOR ALLERGENS  
FROM AMBROSIA ARTEMISIIFOLIA

15

(iii) NUMBER OF SEQUENCES: 150

(iv) CORRESPONDENCE ADDRESS:

20

(A) ADDRESSEE: IMMULOGIC PHARAMCEUTICAL CORPORATION  
(B) STREET: 610 Lincoln Street  
(C) CITY: Waltham  
(D) STATE: MA  
(E) COUNTRY: USA  
(F) ZIP: 02154

25

(v) COMPUTER READABLE FORM:

30

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: ASCII Text

(vi) CURRENT APPLICATION DATA:

35

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

40

(A) APPLICATION NUMBER: US08/298,542  
(B) FILING DATE: August 30, 1994

(viii) ATTORNEY/AGENT INFORMATION:

45

(A) NAME: Darlene A. Vanstone  
(B) REGISTRATION NUMBER: 35,729  
(C) REFERENCE/DOCKET NUMBER: 001.4 US

(ix) TELECOMMUNICATION INFORMATION:

50

(A) TELEPHONE: (617) 466-6000  
(B) TELEFAX: (617) 466-6040

(2) INFORMATION FOR SEQ ID NO:1:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1223 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA



- 55 -

## (ix) FEATURE:

(A) NAME/KEY: CDS

5 (B) LOCATION: 1..1188

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10	ATG GGG ATC AAA CAC TGT TGT TAC ATC TTG TAT TTT ACC TTA GCC CTT	48
	Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu	
	1 5 10 15	
15	GTC ACT TTG CTG CAA CCT GTT CGT TCT GCC GAA GAT CTC CAG GAA ATC	96
	Val Thr Leu Leu Gln Pro Val Arg Ser Ala Glu Asp Leu Gln Glu Ile	
	20 25 30	
20	TTA CCA GTT AAC GAA ACA AGG AGG CTG ACA ACA AGT GGA GCA TAC AAC	144
	Leu Pro Val Asn Glu Thr Arg Arg Leu Thr Thr Ser Gly Ala Tyr Asn	
	35 40 45	
25	ATT ATA GAC GGG TGC TGG AGG GGC AAA GCC GAT TGG GCG GAA AAC CGA	192
	Ile Ile Asp Gly Cys Trp Arg Gly Lys Ala Asp Trp Ala Glu Asn Arg	
	50 55 60	
30	AAA GCG TTA GCC GAT TGT GCC CAA GGT TTT GGG AAG GGA ACA GTG GGC	240
	Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys Gly Thr Val Gly	
	65 70 75 80	
35	GGA AAA GAT GGT GAT ATA TAC ACG GTC ACC AGT GAG CTA GAT GAT GAT	288
	Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp	
	85 90 95	
40	GTT GCA AAT CCA AAA GAA GGC ACA CTC CGG TTT GGT GCC GCC CAA AAC	336
	Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Ala Gln Asn	
	100 105 110	
45	AGG CCC TTG TGG ATC ATT TTT GAA AGA GAT ATG GTG ATT CGT TTG GAT	384
	Arg Pro Leu Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu Asp	
	115 120 125	
50	AAA GAG ATG GTG GTA AAC AGT GAC AAG ACC ATC GAT GGC CGA GGG GCG	432
	Lys Glu Met Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg Gly Ala	
	130 135 140	
55	AAA GTT GAA ATC ATT AAC GCT GGT TTC ACC CTT AAT GGT GTC AAG AAT	480
	Lys Val Glu Ile Ile Asn Ala Gly Phe Thr Leu Asn Gly Val Lys Asn	
	145 150 155 160	
60	GTA ATC ATT CAT AAC ATA AAT ATG CAT GAT GTT AAA GTG AAT CCA GGA	528
	Val Ile Ile His Asn Ile Asn Met His Asp Val Lys Val Asn Pro Gly	
	165 170 175	
65	GGC CTG ATT AAG TCC AAC GAT GGT CCA GCA GCT CCA AGA GCT GGT AGT	576
	Gly Leu Ile Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser	
	180 185 190	
70	GAT GGT GAT GCT ATA AGT ATT TCT GGT AGT TCA CAA ATA TGG ATC GAC	624
	Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp	

- 56 -

	195	200	205	
5	CAT TGT TCG CTC AGT AAG TCT GTT GAT GGG CTG GTA GAT GCC AAG CTC His Cys Ser Leu Ser Lys Ser Val Asp Gly Leu Val Asp Ala Lys Leu 210 215 220			672
10	GGC ACC ACA CGC TTA ACC GTT TCC AAC AGC TTA TTC ACC CAA CAC CAG Gly Thr Thr Arg Leu Thr Val Ser Asn Ser Leu Phe Thr Gln His Gln 225 230 235 240			720
15	TTT GTA CTA TTA TTC GGG GCT GGT GAC GAA AAT ATT GAA GAT AGA GGC Phe Val Leu Leu Phe Gly Ala Gly Asp Glu Asn Ile Glu Asp Arg Gly 245 250 255			768
20	ATG CTA GCA ACG GTC GCT TTC AAC ACG TTC ACT GAT AAC GTT GAC CAA Met Leu Ala Thr Val Ala Phe Asn Thr Phe Thr Asp Asn Val Asp Gln 260 265 270			816
25	AGA ATG CCT AGA TGT CGA CAT GGG TTT TTC CAA GTC GTT AAC AAC AAC Arg Met Pro Arg Cys Arg His Gly Phe Phe Gln Val Val Asn Asn Asn 275 280 285			864
30	TAT GAT AAA TGG GGA TCG TAT GCC ATC GGT GGT AGC GCG TCC CCA ACC Tyr Asp Lys Trp Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr 290 295 300			912
35	ATA CTC AGC CAA GGG AAC AGA TTC TGC GCC CCC GAT GAA CGC AGC AAG Ile Leu Ser Gln Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys 305 310 315 320			960
40	AAA AAT GTC CTA GGA AGG CAT GGT GAA GCC GCC GCA GAG TCG ATG AAG Lys Asn Val Leu Gly Arg His Gly Glu Ala Ala Ala Glu Ser Met Lys 325 330 335			1008
45	TGG AAC TGG AGA ACG AAT AAA GAC GTG CTT GAA AAT GGT GCT ATT TTT Trp Asn Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile Phe 340 345 350			1056
50	GTT GCA TCC GGG GTC GAT CCA GTG CTA ACC CCT GAG CAA AGC GCA GGG Val Ala Ser Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala Gly 355 360 365			1104
55	ATG ATT CCA GCC GAA CCA GGA GAG TCC GCT CTA AGC CTC ACT AGT AGT Met Ile Pro Ala Glu Pro Gly Glu Ser Ala Leu Ser Leu Thr Ser Ser 370 375 380			1152
60	GCT GGT GTA CTC TCA TGC CAA CCC GGA GCA CCT TGC TAAGCACCCG Ala Gly Val Leu Ser Cys Gln Pro Gly Ala Pro Cys 385 390 395			1198
65	ACCAATTACT AAGCACTTAT AATGA			1223

(2) INFORMATION FOR SEQ ID NO:2:

55

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 396 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 57 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

5 Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
    1                  5                  10                  15  
 10 Val Thr Leu Leu Gln Pro Val Arg Ser Ala Glu Asp Leu Gln Glu Ile  
                   20                  25                  30  
   Leu Pro Val Asn Glu Thr Arg Arg Leu Thr Thr Ser Gly Ala Tyr Asn  
           35                  40                  45  
 15 Ile Ile Asp Gly Cys Trp Arg Gly Lys Ala Asp Trp Ala Glu Asn Arg  
       50                  55                  60  
   Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys Gly Thr Val Gly  
       65                  70                  75                  80  
 20 Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp  
                   85                  90                  95  
   Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Ala Gln Asn  
 25                  100                  105                  110  
   Arg Pro Leu Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu Asp  
           115                  120                  125  
 30 Lys Glu Met Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg Gly Ala  
       130                  135                  140  
   Lys Val Glu Ile Ile Asn Ala Gly Phe Thr Leu Asn Gly Val Lys Asn  
 35      145                  150                  155                  160  
   Val Ile Ile His Asn Ile Asn Met His Asp Val Lys Val Asn Pro Gly  
           165                  170                  175  
   Gly Leu Ile Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser  
 40                  180                  185                  190  
   Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp  
       195                  200                  205  
 45 His Cys Ser Leu Ser Lys Ser Val Asp Gly Leu Val Asp Ala Lys Leu  
       210                  215                  220  
   Gly Thr Thr Arg Leu Thr Val Ser Asn Ser Leu Phe Thr Gln His Gln  
       225                  230                  235                  240  
 50 Phe Val Leu Leu Phe Gly Ala Gly Asp Glu Asn Ile Glu Asp Arg Gly  
           245                  250                  255  
   Met Leu Ala Thr Val Ala Phe Asn Thr Phe Thr Asp Asn Val Asp Gln  
 55          260                  265                  270  
   Arg Met Pro Arg Cys Arg His Gly Phe Phe Gln Val Val Asn Asn Asn  
       275                  280                  285

- 58 -

Tyr Asp Lys Trp Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr  
 290 295 300  
 Ile Leu Ser Gln Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys  
 5 305 310 315 320  
 Lys Asn Val Leu Gly Arg His Gly Glu Ala Ala Ala Glu Ser Met Lys  
 325 330 335  
 10 Trp Asn Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile Phe  
 340 345 350  
 Val Ala Ser Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala Gly  
 355 360 365  
 15 Met Ile Pro Ala Glu Pro Gly Glu Ser Ala Leu Ser Leu Thr Ser Ser  
 370 375 380  
 Ala Gly Val Leu Ser Cys Gln Pro Gly Ala Pro Cys  
 20 385 390 395

## (2) INFORMATION FOR SEQ ID NO:3:

25 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1349 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

35 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1194

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

40 ATG GGG ATC AAA CAC TGT TGT TAC ATC TTG TAT TTT ACC TTA GCC CTT 48  
 Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
 1 5 10 15  
 45 GTC ACT TTG CTG CAA CCT GTT CGT TCT GCA GAA GAT GTT GAA GAA TTC 96  
 Val Thr Leu Leu Gln Pro Val Arg Ser Ala Glu Asp Val Glu Glu Phe  
 20 25 30  
 50 TTA CCT TCA GCT AAC GAA ACA AGG AGG AGC CTG AAA GCA TGT GAA GCA 144  
 Leu Pro Ser Ala Asn Glu Thr Arg Arg Ser Leu Lys Ala Cys Glu Ala  
 35 40 45  
 CAC AAC ATT ATA GAC AAG TGC TGG AGG TGC AAA GCC GAT TGG GCG AAT 192  
 His Asn Ile Ile Asp Lys Cys Trp Arg Cys Lys Ala Asp Trp Ala Asn  
 55 50 55 60  
 AAC CGA CAA GCG TTA GCC GAT TGT GCC CAA GGT TTT GCA AAG GGA ACC 240  
 Asn Arg Gln Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys Gly Thr  
 65 70 75 80

	TAC GGT GGA AAA CAT GGT GAT GTC TAC ACG GTC ACC AGT GAT AAA GAT	288
	Tyr Gly Gly Lys His Gly Asp Val Tyr Thr Val Thr Ser Asp Lys Asp	
	85 90 95	
5	GAT GAT GTT GCA AAT CCA AAA GAA GGC ACA CTC CGG TTT GCT GCT GCC	336
	Asp Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Ala Ala Ala	
	100 105 110	
10	CAA AAC AGG CCC TTG TGG ATC ATT TTT AAA AGA AAT ATG GTG ATT CAT	384
	Gln Asn Arg Pro Leu Trp Ile Ile Phe Lys Arg Asn Met Val Ile His	
	115 120 125	
15	TTG AAT CAA GAG CTT GTC GTA AAC AGC GAC AAG ACC ATC GAT GGC CGA	432
	Leu Asn Gln Glu Leu Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg	
	130 135 140	
20	GGG GTG AAA GTT AAC ATC GTT AAC GCC GGT CTC ACC CTC ATG AAT GTC	480
	Gly Val Lys Val Asn Ile Val Asn Ala Gly Leu Thr Leu Met Asn Val	
	145 150 155 160	
25	AAG AAT ATA ATC ATT CAT AAC ATA AAT ATC CAT GAT ATT AAA GTT TGT	528
	Lys Asn Ile Ile Ile His Asn Ile Asn Ile His Asp Ile Lys Val Cys	
	165 170 175	
30	CCA GGA GGC ATG ATT AAG TCC AAC GAT GGT CCA CCA ATT TTA AGA CAA	576
	Pro Gly Gly Met Ile Lys Ser Asn Asp Gly Pro Pro Ile Leu Arg Gln	
	180 185 190	
35	CAA AGT GAT GGT GAT GCT ATA AAT GTT GCT GGT AGT TCA CAA ATA TGG	624
	Gln Ser Asp Gly Asp Ala Ile Asn Val Ala Gly Ser Ser Gln Ile Trp	
	195 200 205	
40	ATC GAC CAT TGC TCG CTC AGT AAG GCT TCC GAT GGG CTG CTC GAT ATC	672
	Ile Asp His Cys Ser Leu Ser Lys Ala Ser Asp Gly Leu Leu Asp Ile	
	210 215 220	
45	ACC CTC GGC AGC TCA CAC GTG ACC GTT TCC AAC TGC AAA TTC ACC CAA	720
	Thr Leu Gly Ser Ser His Val Thr Val Ser Asn Cys Lys Phe Thr Gln	
	225 230 235 240	
50	CAC CAA TTT GTA TTA TTG CTC GGG GCT GAT GAC ACC CAT TAT CAA GAT	768
	His Gln Phe Val Leu Leu Leu Gly Ala Asp Asp Thr His Tyr Gln Asp	
	245 250 255	
55	AAA GGC ATG CTA GCA ACG GTA GCA TTC AAC ATG TTC ACC GAT CAC GTT	816
	Lys Gly Met Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp His Val	
	260 265 270	
60	GAC CAA AGA ATG CCT AGA TGT AGA TTT GGG TTT TTC CAA GTC GTT AAC	864
	Asp Gln Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Val Val Asn	
	275 280 285	
65	AAC AAC TAC GAC AGA TGG GGA ACG TAC GCC ATC GGT GGT AGC TCG GCC	912
	Asn Asn Tyr Asp Arg Trp Gly Thr Tyr Ala Ile Gly Gly Ser Ser Ala	
	290 295 300	
70	CCA ACT ATA CTC AGC CAA GGG AAC AGA TTC TTC GCC CCC GAT GAT ATC	960
	Pro Thr Ile Leu Ser Gln Gly Asn Arg Phe Phe Ala Pro Asp Asp Ile	

- 60 -

305                      310                      315                      320  
 ATC AAG AAA AAT GTC TTA GCG AGG ACT GGT ACT GGC AAC GCA GAG TCG      1008  
 Ile Lys Lys Asn Val Leu Ala Arg Thr Gly Thr Gly Asn Ala Glu Ser  
 5                      325                      330                      335  
 ATG TCG TGG AAC TGG AGA ACA GAT AGA GAC TTG CTT GAA AAT GGT GCT      1056  
 Met Ser Trp Asn Trp Arg Thr Asp Arg Asp Leu Leu Glu Asn Gly Ala  
 340                      345                      350  
 10      ATT TTT CTC CCA TCC GGG TCT GAT CCA GTG CTA ACC CCT GAG CAA AAA      1104  
 Ile Phe Leu Pro Ser Gly Ser Asp Pro Val Leu Thr Pro Glu Gln Lys  
 355                      360                      365  
 15      GCA GGG ATG ATT CCA GCT GAA CCA GGA GAA GCC GTT CTA AGA CTC ACT      1152  
 Ala Gly Met Ile Pro Ala Glu Pro Gly Glu Ala Val Leu Arg Leu Thr  
 370                      375                      380  
 20      AGT AGT GCT GGT GTA CTC TCA TGC CAT CAA GGA GCA CCT TGC TAA      1197  
 Ser Ser Ala Gly Val Leu Ser Cys His Gln Gly Ala Pro Cys  
 385                      390                      395  
 GCACCTGGCC AATTCCTAAG CTTTATAAT AATCATAAAT ACTTATTTTA TTTTATTTTT      1260  
 25      GATATTTTAT ATGAACCATT ACGTTCAAGT ACTCTATTAA CATGTTTTAA ATTCATAAGA      1320  
 GTTTATTGAT AAAAAAAAAA AAAACCGAAT TC      1349

30      (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 398 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

40      Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
          1                      5                      10                      15  
 45      Val Thr Leu Leu Gln Pro Val Arg Ser Ala Glu Asp Val Glu Glu Phe  
          20                      25                      30  
 Leu Pro Ser Ala Asn Glu Thr Arg Arg Ser Leu Lys Ala Cys Glu Ala  
          35                      40                      45  
 50      His Asn Ile Ile Asp Lys Cys Trp Arg Cys Lys Ala Asp Trp Ala Asn  
          50                      55                      60  
 Asn Arg Gln Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys Gly Thr  
          65                      70                      75                      80  
 55      Tyr Gly Gly Lys His Gly Asp Val Tyr Thr Val Thr Ser Asp Lys Asp  
          85                      90                      95  
 Asp Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Ala Ala Ala

- 61 -

	100	105	110
	Gln Asn Arg Pro Leu Trp Ile Ile Phe Lys Arg Asn Met Val Ile His		
	115	120	125
5	Leu Asn Gln Glu Leu Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg		
	130	135	140
	Gly Val Lys Val Asn Ile Val Asn Ala Gly Leu Thr Leu Met Asn Val		
10	145	150	155 160
	Lys Asn Ile Ile Ile His Asn Ile Asn Ile His Asp Ile Lys Val Cys		
	165	170	175
15	Pro Gly Gly Met Ile Lys Ser Asn Asp Gly Pro Pro Ile Leu Arg Gln		
	180	185	190
	Gln Ser Asp Gly Asp Ala Ile Asn Val Ala Gly Ser Ser Gln Ile Trp		
20	195	200	205
	Ile Asp His Cys Ser Leu Ser Lys Ala Ser Asp Gly Leu Leu Asp Ile		
	210	215	220
	Thr Leu Gly Ser Ser His Val Thr Val Ser Asn Cys Lys Phe Thr Gln		
25	225	230	235 240
	His Gln Phe Val Leu Leu Leu Gly Ala Asp Asp Thr His Tyr Gln Asp		
	245	250	255
30	Lys Gly Met Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp His Val		
	260	265	270
	Asp Gln Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Val Val Asn		
35	275	280	285
	Asn Asn Tyr Asp Arg Trp Gly Thr Tyr Ala Ile Gly Gly Ser Ser Ala		
	290	295	300
	Pro Thr Ile Leu Ser Gln Gly Asn Arg Phe Phe Ala Pro Asp Asp Ile		
40	305	310	315 320
	Ile Lys Lys Asn Val Leu Ala Arg Thr Gly Thr Gly Asn Ala Glu Ser		
	325	330	335
45	Met Ser Trp Asn Trp Arg Thr Asp Arg Asp Leu Leu Glu Asn Gly Ala		
	340	345	350
	Ile Phe Leu Pro Ser Gly Ser Asp Pro Val Leu Thr Pro Glu Gln Lys		
50	355	360	365
	Ala Gly Met Ile Pro Ala Glu Pro Gly Glu Ala Val Leu Arg Leu Thr		
	370	375	380
	Ser Ser Ala Gly Val Leu Ser Cys His Gln Gly Ala Pro Cys		
55	385	390	395

- 62 -

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1320 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

5

## (ii) MOLECULE TYPE: cDNA

10

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1191

15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

1	ATG GGG ATC AAA CAA TGT TGT TAC ATC TTG TAT TTT ACC TTA GCA CTT	48
	Met Gly Ile Lys Gln Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu	
5		15
20	GTC GCT TTG CTG CAA CCT GTT CGT TCT GCC GAA GGT GTC GGG GAA ATC	96
	Val Ala Leu Leu Gln Pro Val Arg Ser Ala Glu Gly Val Gly Glu Ile	
25		30
35	TTA CCT TCA GTT AAC GAA ACG AGG AGC CTG CAA GCA TGT GAA GCA CTC	144
	Leu Pro Ser Val Asn Glu Thr Arg Ser Leu Gln Ala Cys Glu Ala Leu	
40		45
50	AAC ATT ATA GAC AAG TGC TGG AGG GGC AAA GCC GAT TGG GAG AAC AAC	192
	Asn Ile Ile Asp Lys Cys Trp Arg Gly Lys Ala Asp Trp Glu Asn Asn	
55		60
65	CGA CAA GCG TTA GCC GAC TGT GCC CAA GGT TTT GCA AAG GGA ACC TAC	240
	Arg Gln Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys Gly Thr Tyr	
70		75
85	GGC GGA AAA TGG GGT GAT GTC TAC ACG GTC ACC AGC AAT CTA GAT GAT	288
	Gly Gly Lys Trp Gly Asp Val Tyr Thr Val Thr Ser Asn Leu Asp Asp	
90		95
100	GAT GTT GCA AAT CCA AAA GAA GGC ACA CTC CGG TTT GCT GCC GCC CAA	336
	Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Ala Ala Gln	
105		110
115	AAC AGG CCC TTG TGG ATC ATT TTT AAA AAT GAT ATG GTG ATT AAT TTG	384
	Asn Arg Pro Leu Trp Ile Ile Phe Lys Asn Asp Met Val Ile Asn Leu	
120		125
130	AAT CAA GAG CTT GTC GTA AAC AGC GAC AAG ACC ATC GAT GGC CGA GGG	432
	Asn Gln Glu Leu Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg Gly	
135		140
145	GTG AAA GTT GAA ATC ATT AAC GGA GGT CTC ACC CTC ATG AAT GTC AAG	480
	Val Lys Val Glu Ile Ile Asn Gly Gly Leu Thr Leu Met Asn Val Lys	
150		155
160	AAT ATA ATC ATT CAT AAC ATA AAT ATC CAT GAT GTT AAA GTG CTT CCA	528
	Asn Ile Ile Ile His Asn Ile Asn Ile His Asp Val Lys Val Leu Pro	
165		170
175		175



- 63 -

	GGA GGC ATG ATT AAG TCC AAC GAT GGT CCA CCA ATT TTA AGA CAA GCA	576
	Gly Gly Met Ile Lys Ser Asn Asp Gly Pro Pro Ile Leu Arg Gln Ala	
	180 185 190	
5	AGT GAT GGG GAT ACT ATA AAT GTT GCT GGT AGT TCC CAA ATA TGG ATA	624
	Ser Asp Gly Asp Thr Ile Asn Val Ala Gly Ser Ser Gln Ile Trp Ile	
	195 200 205	
10	GAC CAT TGC TCA CTC AGC AAG TCT TTC GAT GGG CTG GTC GAT GTC ACC	672
	Asp His Cys Ser Leu Ser Lys Ser Phe Asp Gly Leu Val Asp Val Thr	
	210 215 220	
15	CTC GGT AGC ACA CAC GTG ACC ATT TCC AAC TGC AAA TTC ACC CAA CAG	720
	Leu Gly Ser Thr His Val Thr Ile Ser Asn Cys Lys Phe Thr Gln Gln	
	225 230 235 240	
20	TCA AAA GCA ATA TTG TTG GGA GCA GAT GAC ACC CAT GTT CAA GAT AAA	768
	Ser Lys Ala Ile Leu Leu Gly Ala Asp Asp Thr His Val Gln Asp Lys	
	245 250 255	
25	GGA ATG CTA GCA ACG GTC GCT TTC AAC ATG TTC ACC GAT AAC GTT GAC	816
	Gly Met Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp Asn Val Asp	
	260 265 270	
30	CAA AGA ATG CCT AGA TGT CGA TTT GGG TTT TTC CAA GTT GTT AAC AAC	864
	Gln Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Val Val Asn Asn	
	275 280 285	
35	AAC TAC GAC AGA TGG GGA ACG TAC GCC ATA GGT GGT AGC TCG GCC CCA	912
	Asn Tyr Asp Arg Trp Gly Thr Tyr Ala Ile Gly Gly Ser Ser Ala Pro	
	290 295 300	
40	ACT ATA CTC TGC CAA GGG AAC AGA TTC TTG GCC CCT GAT GAT CAG ATC	960
	Thr Ile Leu Cys Gln Gly Asn Arg Phe Leu Ala Pro Asp Asp Gln Ile	
	305 310 315 320	
45	AAG AAA AAT GTC CTA GCG AGG ACT GGT ACA GGC GCT GCT GAG TCG ATG	1008
	Lys Lys Asn Val Leu Ala Arg Thr Gly Thr Gly Ala Ala Glu Ser Met	
	325 330 335	
50	GCG TGG AAC TGG AGA TCT GAT AAA GAC TTG CTT GAA AAT GGT GCT ATT	1056
	Ala Trp Asn Trp Arg Ser Asp Lys Asp Leu Leu Glu Asn Gly Ala Ile	
	340 345 350	
55	TTT GTT ACA TCT GGG TCT GAT CCA GTG CTA ACC CCT GTT CAA AGC GCA	1104
	Phe Val Thr Ser Gly Ser Asp Pro Val Leu Thr Pro Val Gln Ser Ala	
	355 360 365	
60	GGG ATG ATT CCA GCT GAA CCA GGA GAA GCC GCT ATA AAA CTC ACT AGT	1152
	Gly Met Ile Pro Ala Glu Pro Gly Glu Ala Ala Ile Lys Leu Thr Ser	
	370 375 380	
65	AGT GCT GGT GTA TTC TCA TGC CGT CCT GGA GCA CCT TGC TAAGCACCT	1201
	Ser Ala Gly Val Phe Ser Cys Arg Pro Gly Ala Pro Cys	
	385 390 395	
70	GCCAATTCTC CTAAGCTTTT GCAATGATCA AAAATACTTT TTTATTTTAT TTTTAATATT	1261

TTATATGTAC TGGAAATGAA CCATTACCTT CTAGTACTCT ATAACATGTT TTGCATTTA

1320

## (2) INFORMATION FOR SEQ ID NO:6:

5

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 397 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

10

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

15 Met Gly Ile Lys Gln Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
     1                    5                    10                    15  
 Val Ala Leu Leu Gln Pro Val Arg Ser Ala Glu Gly Val Gly Glu Ile  
                     20                    25                    30  
 20 Leu Pro Ser Val Asn Glu Thr Arg Ser Leu Gln Ala Cys Glu Ala Leu  
                     35                    40                    45  
 Asn Ile Ile Asp Lys Cys Trp Arg Gly Lys Ala Asp Trp Glu Asn Asn  
 25                    50                    55                    60  
 Arg Gln Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys Gly Thr Tyr  
                     65                    70                    75                    80  
 30 Gly Gly Lys Trp Gly Asp Val Tyr Thr Val Thr Ser Asn Leu Asp Asp  
                     85                    90                    95  
 Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Ala Ala Ala Gln  
                     100                    105                    110  
 35 Asn Arg Pro Leu Trp Ile Ile Phe Lys Asn Asp Met Val Ile Asn Leu  
                     115                    120                    125  
 Asn Gln Glu Leu Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg Gly  
 40                    130                    135                    140  
 Val Lys Val Glu Ile Ile Asn Gly Gly Leu Thr Leu Met Asn Val Lys  
                     145                    150                    155                    160  
 45 Asn Ile Ile Ile His Asn Ile Asn Ile His Asp Val Lys Val Leu Pro  
                     165                    170                    175  
 Gly Gly Met Ile Lys Ser Asn Asp Gly Pro Pro Ile Leu Arg Gln Ala  
                     180                    185                    190  
 50 Ser Asp Gly Asp Thr Ile Asn Val Ala Gly Ser Ser Gln Ile Trp Ile  
                     195                    200                    205  
 Asp His Cys Ser Leu Ser Lys Ser Phe Asp Gly Leu Val Asp Val Thr  
 55                    210                    215                    220  
 Leu Gly Ser Thr His Val Thr Ile Ser Asn Cys Lys Phe Thr Gln Gln  
                     225                    230                    235                    240

- 65 -

Ser Lys Ala Ile Leu Leu Gly Ala Asp Asp Thr His Val Gln Asp Lys  
 245 250 255

5 Gly Met Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp Asn Val Asp  
 260 265 270

Gln Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Val Val Asn Asn  
 275 280 285

10 Asn Tyr Asp Arg Trp Gly Thr Tyr Ala Ile Gly Gly Ser Ser Ala Pro  
 290 295 300

Thr Ile Leu Cys Gln Gly Asn Arg Phe Leu Ala Pro Asp Asp Gln Ile  
 305 310 315 320

15 Lys Lys Asn Val Leu Ala Arg Thr Gly Thr Gly Ala Ala Glu Ser Met  
 325 330 335

20 Ala Trp Asn Trp Arg Ser Asp Lys Asp Leu Leu Glu Asn Gly Ala Ile  
 340 345 350

Phe Val Thr Ser Gly Ser Asp Pro Val Leu Thr Pro Val Gln Ser Ala  
 355 360 365

25 Gly Met Ile Pro Ala Glu Pro Gly Glu Ala Ala Ile Lys Leu Thr Ser  
 370 375 380

Ser Ala Gly Val Phe Ser Cys Arg Pro Gly Ala Pro Cys  
 385 390 395

30

## (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:  
 35 (A) LENGTH: 1187 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:  
 45 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1176

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

50 ATG GGG ATC AAA CAC TGT TGT TAC ATC TTG TAT TTT ACC TTA GCC CTT 48  
 Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
 1 5 10 15

GTC ACT TTG CTG CAA CCT GTT CGT TCT GCC GAA GAT CTC CAG GAA ATC 96  
 55 Val Thr Leu Leu Gln Pro Val Arg Ser Ala Glu Asp Leu Gln Glu Ile  
 20 25 30

TTA CCT TCA GCT AAC GAA ACA AGG AGC CTG ACA ACA TGT GGA ACA TAC 144  
 Leu Pro Ser Ala Asn Glu Thr Arg Ser Leu Thr Thr Cys Gly Thr Tyr

- 66 -

	35	40	45	
5	AAC ATT ATA GAC GGG TGC TGG AGG GGC AAA GCC GAT TGG GCG GAA AAC Asn Ile Ile Asp Gly Cys Trp Arg Gly Lys Ala Asp Trp Ala Glu Asn 50 55 60			192
10	CGA AAA GCG TTA GCC GAT TGT GCC CAA GGT TTT GCA AAG GGA ACA ATC Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys Gly Thr Ile 65 70 75 80			240
15	GGC GGA AAA GAT GGT GAT ATA TAC ACG GTC ACC AGT GAG CTA GAT GAT Gly Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp 85 90 95			288
20	GAT GTT GCA AAT CCA AAA GAA GGC ACA CTC CGG TTT GGT GCC GCC CAA Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Ala Gln 100 105 110			336
25	AAC AGG CCC TTG TGG ATT ATT TTT GAA AGA GAT ATG GTG ATT CGT TTG Asn Arg Pro Leu Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu 115 120 125			384
30	GAT AGA GAG TTG GCT ATA AAC AAC GAC AAG ACC ATC GAT GGC CGA GGG Asp Arg Glu Leu Ala Ile Asn Asn Asp Lys Thr Ile Asp Gly Arg Gly 130 135 140			432
35	GCG AAA GTT GAA ATC ATT AAC GCT GGT TTC GCC ATC TAT AAT GTC AAG Ala Lys Val Glu Ile Ile Asn Ala Gly Phe Ala Ile Tyr Asn Val Lys 145 150 155 160			480
40	AAT ATA ATC ATT CAT AAC ATA ATT ATG CAT GAT ATT GTA GTG AAT CCA Asn Ile Ile Ile His Asn Ile Ile Met His Asp Ile Val Val Asn Pro 165 170 175			528
45	GGA GGC CTG ATT AAG TCC CAC GAT GGT CCA CCA GTT CCA AGA AAG GGT Gly Gly Leu Ile Lys Ser His Asp Gly Pro Pro Val Pro Arg Lys Gly 180 185 190			576
50	AGT GAT GGT GAT GCT ATA GGT ATT TCT GGT GGT TCA CAA ATA TGG ATC Ser Asp Gly Asp Ala Ile Gly Ile Ser Gly Gly Ser Gln Ile Trp Ile 195 200 205			624
55	GAC CAT TGC TCC CTC AGT AAG GCT GTT GAT GGG CTA ATC GAT GCT AAA Asp His Cys Ser Leu Ser Lys Ala Val Asp Gly Leu Ile Asp Ala Lys 210 215 220			672
60	CAC GGC AGC ACA CAC TTC ACC GTT TCT AAC TGC TTA TTC ACC CAA CAC His Gly Ser Thr His Phe Thr Val Ser Asn/Cys Leu Phe Thr Gln His 225 230 235 240			720
65	CAA TAT TTA TTA TTG TTC TGG GAT TTT GAC GAG CGA GGC ATG CTA TGT Gln Tyr Leu Leu Leu Phe Trp Asp Phe Asp Glu Arg Gly Met Leu Cys 245 250 255			768
70	ACG GTC GCA TTC AAC AAG TTC ACT GAT AAC GTT GAC CAA AGA ATG CCT Thr Val Ala Phe Asn Lys Phe Thr Asp Asn Val Asp Gln Arg Met Pro 260 265 270			816
75	AAC TTA CGA CAT GGG TTT GTC CAA GTC GTT AAC AAC AAC TAC GAA AGA			864

- 67 -

	Asn	Leu	Arg	His	Gly	Phe	Val	Gln	Val	Val	Asn	Asn	Asn	Tyr	Glu	Arg	
		275						280					285				
5	TGG	GGA	TCG	TAC	GCC	CTC	GGT	GGT	AGC	GCA	GGC	CCA	ACC	ATA	CTT	AGC	912
	Trp	Gly	Ser	Tyr	Ala	Leu	Gly	Gly	Ser	Ala	Gly	Pro	Thr	Ile	Leu	Ser	
		290					295					300					
10	CAA	GGG	AAC	AGA	TTC	TTA	GCC	TCC	GAT	ATC	AAG	AAA	GAG	GTC	GTA	GGG	960
	Gln	Gly	Asn	Arg	Phe	Leu	Ala	Ser	Asp	Ile	Lys	Lys	Glu	Val	Val	Gly	
		305				310					315					320	
15	AGG	TAT	GGT	GAA	TCC	GCC	ATG	TCA	GAG	TCG	ATT	AAT	TGG	AAC	TGG	AGA	1008
	Arg	Tyr	Gly	Glu	Ser	Ala	Met	Ser	Glu	Ser	Ile	Asn	Trp	Asn	Trp	Arg	
	—				325					330					335		
20	TCG	TAT	ATG	GAC	GTA	TTT	GAA	AAT	GGT	GCT	ATT	TTT	GTT	CCA	TCC	GGG	1056
	Ser	Tyr	Met	Asp	Val	Phe	Glu	Asn	Gly	Ala	Ile	Phe	Val	Pro	Ser	Gly	
				340					345					350			
25	GTT	GAT	CCA	GTG	CTA	ACC	CCT	GAG	CAA	AAC	GCA	GGG	ATG	ATT	CCA	GCC	1104
	Val	Asp	Pro	Val	Leu	Thr	Pro	Glu	Gln	Asn	Ala	Gly	Met	Ile	Pro	Ala	
		355					360					365					
30	GAA	CCA	GGA	GAA	GCC	GTT	CTA	AGA	CTC	ACT	AGT	AGT	GCT	GGT	GTC	CTC	1152
	Glu	Pro	Gly	Glu	Ala	Val	Leu	Arg	Leu	Thr	Ser	Ser	Ala	Gly	Val	Leu	
		370				375					380						
35	TCA	TGC	CAA	CCT	GGA	GCA	CCT	TGC	TAAGCACTGC	A							1187
	Ser	Cys	Gln	Pro	Gly	Ala	Pro	Cys									
		385				390											

## (2) INFORMATION FOR SEQ ID NO:8:

35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 392 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

45	Met	Gly	Ile	Lys	His	Cys	Cys	Tyr	Ile	Leu	Tyr	Phe	Thr	Leu	Ala	Leu	
	1				5					10					15		
	Val	Thr	Leu	Leu	Gln	Pro	Val	Arg	Ser	Ala	Glu	Asp	Leu	Gln	Glu	Ile	
			20					25					30				
50	Leu	Pro	Ser	Ala	Asn	Glu	Thr	Arg	Ser	Leu	Thr	Thr	Cys	Gly	Thr	Tyr	
			35					40					45				
	Asn	Ile	Ile	Asp	Gly	Cys	Trp	Arg	Gly	Lys	Ala	Asp	Trp	Ala	Glu	Asn	
		50				55					60						
55	Arg	Lys	Ala	Leu	Ala	Asp	Cys	Ala	Gln	Gly	Phe	Ala	Lys	Gly	Thr	Ile	
		65				70					75					80	
	Gly	Gly	Lys	Asp	Gly	Asp	Ile	Tyr	Thr	Val	Thr	Ser	Glu	Leu	Asp	Asp	

- 68 -

	85	90	95
	Asp Val Ala Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Ala Gln		
	100	105	110
5	Asn Arg Pro Leu Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu		
	115	120	125
10	Asp Arg Glu Leu Ala Ile Asn Asn Asp Lys Thr Ile Asp Gly Arg Gly		
	130	135	140
	Ala Lys Val Glu Ile Ile Asn Ala Gly Phe Ala Ile Tyr Asn Val Lys		
	145	150	155
	150		160
15	Asn Ile Ile Ile His Asn Ile Ile Met His Asp Ile Val Val Asn Pro		
	165	170	175
	Gly Gly Leu Ile Lys Ser His Asp Gly Pro Pro Val Pro Arg Lys Gly		
	180	185	190
20	Ser Asp Gly Asp Ala Ile Gly Ile Ser Gly Gly Ser Gln Ile Trp Ile		
	195	200	205
	Asp His Cys Ser Leu Ser Lys Ala Val Asp Gly Leu Ile Asp Ala Lys		
25	210	215	220
	His Gly Ser Thr His Phe Thr Val Ser Asn Cys Leu Phe Thr Gln His		
	225	230	235
			240
30	Gln Tyr Leu Leu Leu Phe Trp Asp Phe Asp Glu Arg Gly Met Leu Cys		
	245	250	255
	Thr Val Ala Phe Asn Lys Phe Thr Asp Asn Val Asp Gln Arg Met Pro		
	260	265	270
35	Asn Leu Arg His Gly Phe Val Gln Val Val Asn Asn Asn Tyr Glu Arg		
	275	280	285
	Trp Gly Ser Tyr Ala Leu Gly Gly Ser Ala Gly Pro Thr Ile Leu Ser		
40	290	295	300
	Gln Gly Asn Arg Phe Leu Ala Ser Asp Ile Lys Lys Glu Val Val Gly		
	305	310	315
			320
45	Arg Tyr Gly Glu Ser Ala Met Ser Glu Ser Ile Asn Trp Asn Trp Arg		
	325	330	335
	Ser Tyr Met Asp Val Phe Glu Asn Gly Ala Ile Phe Val Pro Ser Gly		
	340	345	350
50	Val Asp Pro Val Leu Thr Pro Glu Gln Asn Ala Gly Met Ile Pro Ala		
	355	360	365
	Glu Pro Gly Glu Ala Val Leu Arg Leu Thr Ser Ser Ala Gly Val Leu		
55	370	375	380
	Ser Cys Gln Pro Gly Ala Pro Cys		
	385	390	

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1395 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## 10 (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- 15 — (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1191

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

20	ATG GGG ATC AAA CAT TGT TGT TAC ATC TTG TAT TTT ACC TTA GCA CTT	48
	Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu	
	1 5 10 15	
25	GTC ACT TTG GTG CAA GCT GGA CGT CTT GGC GAA GAG GTC GAC ATC TTA	96
	Val Thr Leu Val Gln Ala Gly Arg Leu Gly Glu Glu Val Asp Ile Leu	
	20 25 30	
30	CCT TCA CCT AAC GAT ACA AGG AGG AGC CTG CAA GGA TGT GAA GCA CAC	144
	Pro Ser Pro Asn Asp Thr Arg Arg Ser Leu Gln Gly Cys Glu Ala His	
	35 40 45	
35	AAC ATT ATA GAC AAG TGT TGG AGG TGC AAA CCC GAT TGG GCG GAG AAC	192
	Asn Ile Ile Asp Lys Cys Trp Arg Cys Lys Pro Asp Trp Ala Glu Asn	
	50 55 60	
40	CGA CAA GCG TTA GGC GAT TGT GCG CAA GGT TTT GGA AAG GCA ACT CAC	240
	Arg Gln Ala Leu Gly Asp Cys Ala Gln Gly Phe Gly Lys Ala Thr His	
	65 70 75 80	
45	GGC GGA AAA TGG GGT GAT ATC TAC ATG GTC ACA AGT GAT CAG GAT GAT	288
	Gly Gly Lys Trp Gly Asp Ile Tyr Met Val Thr Ser Asp Gln Asp Asp	
	85 90 95	
50	GAT GTT GTA AAT CCA AAA GAA GGC ACA CTC CGG TTC GGT GCT ACC CAG	336
	Asp Val Val Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Thr Gln	
	100 105 110	
55	GAC AGG CCC TTG TGG ATC ATT TTT CAA AGA GAT ATG ATT ATT TAT TTG	384
	Asp Arg Pro Leu Trp Ile Ile Phe Gln Arg Asp Met Ile Ile Tyr Leu	
	115 120 125	
60	CAA CAA GAG ATG GTC GTA ACC AGC GAC ACC ACC ATT GAT GGT CGA GGG	432
	Gln Gln Glu Met Val Val Thr Ser Asp Thr Thr Ile Asp Gly Arg Gly	
	130 135 140	
65	GCG AAA GTT GAG CTC GTT TAT GGA GGT ATC ACC CTC ATG AAT GTC AAG	480
	Ala Lys Val Glu Leu Val Tyr Gly Gly Ile Thr Leu Met Asn Val Lys	
	145 150 155 160	

- 70 -

	AAT GTA ATC ATT CAC AAC ATA GAT ATC CAT GAT GTT AGA GTG CTT CCA Asn Val Ile Ile His Asn Ile Asp Ile His Asp Val Arg Val Leu Pro	528
	165 170 175	
5	GGA GGT AGG ATT AAG TCC AAT GGT GGT CCA GCC ATA CCA AGA CAT CAG Gly Gly Arg Ile Lys Ser Asn Gly Gly Pro Ala Ile Pro Arg His Gln	576
	180 185 190	
10	AGT GAT GGT GAT GCT ATC CAT GTT ACG GGT AGT TCA GAC ATA TGG ATC Ser Asp Gly Asp Ala Ile His Val Thr Gly Ser Ser Asp Ile Trp Ile	624
	195 200 205	
15	GAC CAT TGC ACG CTC AGT AAG TCA TTT GAT GGG CTC GTC GAT GTC AAC Asp His Cys Thr Leu Ser Lys Ser Phe Asp Gly Leu Val Asp Val Asn	672
	210 215 220	
20	TGG GGC AGC ACA GGA GTA ACC ATT TCC AAC TGC AAA TTC ACC CAC CAC Trp Gly Ser Thr Gly Val Thr Ile Ser Asn Cys Lys Phe Thr His His	720
	225 230 235 240	
25	GAA AAA GCT GTT TTG CTC GGG GCT AGT GAC ACG CAT TTT CAA GAT CTG Glu Lys Ala Val Leu Leu Gly Ala Ser Asp Thr His Phe Gln Asp Leu	768
	245 250 255	
30	AAA ATG CAT GTA ACG CTT GCA TAC AAC ATC TTC ACC AAT ACC GTT CAC Lys Met His Val Thr Leu Ala Tyr Asn Ile Phe Thr Asn Thr Val His	816
	260 265 270	
35	GAA AGA ATG CCC AGA TGC CGA TTT GGG TTT TTC CAA ATC GTT AAC AAC Glu Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Ile Val Asn Asn	864
	275 280 285	
40	TTC TAC GAC AGA TGG GAT AAG TAC GCC ATC GGT GGT AGC TCG AAC CCT Phe Tyr Asp Arg Trp Asp Lys Tyr Ala Ile Gly Gly Ser Ser Asn Pro	912
	290 295 300	
45	ACT ATT CTC AGC CAA GGG AAC AAA TTC GTG GCC CCC GAT TTC ATT TAC Thr Ile Leu Ser Gln Gly Asn Lys Phe Val Ala Pro Asp Phe Ile Tyr	960
	305 310 315 320	
50	AAG AAA AAC GTC TGT CTA AGG ACT GGT GCA CAG GAG CCA GAA TGG ATG Lys Lys Asn Val Cys Leu Arg Thr Gly Ala Gln Glu Pro Glu Trp Met	1008
	325 330 335	
55	ACT TGG AAC TGG AGA ACA CAA AAC GAC GTG CTT GAA AAT GGT GCT ATC Thr Trp Asn Trp Arg Thr Gln Asn Asp Val Leu Glu Asn Gly Ala Ile	1056
	340 345 350	
60	TTT GTG GCA TCT GGG TCT GAT CCA GTG CTA ACC GCT GAA CAA AAT GCA Phe Val Ala Ser Gly Ser Asp Pro Val Leu Thr Ala Glu Gln Asn Ala	1104
	355 360 365	
65	GGC ATG ATG CAA GCT GAA CCG GGA GAT ATG GTT CCA CAA CTC ACC ATG Gly Met Met Gln Ala Glu Pro Gly Asp Met Val Pro Gln Leu Thr Met	1152
	370 375 380	
70	AAT GCA GGT GTA CTC ACA TGC TCG CCT GGA GCA CCT TGC TAAGCACCTG Asn Ala Gly Val Leu Thr Cys Ser Pro Gly Ala Pro Cys	1201
	385 390 395	



5      GCCAATTCCT ATGCAACGAT CATAAATACT TGCTCACCAT AAGTGTTTCAT TTGATTAGAT      1261  
       TTGGACACGA ATGATGTAAC CGATTCTGCT GAATTATGAT TTGTTTTGAT TCTCAGTTTC      1321  
       ATAATATGGC TTCTTGAGAG CAAAATTAGA GAAGAGTGTC TTTGATCAAC TACATTTTAT      1381  
       GGTTTTTATA TTAA      1395

10

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

15      -      (A) LENGTH: 397 amino acids  
               (B) TYPE: amino acid  
               (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Gly Ile Lys His Cys Cys Tyr Ile Leu Tyr Phe Thr Leu Ala Leu  
   1                  5                  10                  15  
 25 Val Thr Leu Val Gln Ala Gly Arg Leu Gly Glu Glu Val Asp Ile Leu  
                   20                  25                  30  
 Pro Ser Pro Asn Asp Thr Arg Arg Ser Leu Gln Gly Cys Glu Ala His  
                   35                  40                  45  
 30 Asn Ile Ile Asp Lys Cys Trp Arg Cys Lys Pro Asp Trp Ala Glu Asn  
                   50                  55                  60  
 35 Arg Gln Ala Leu Gly Asp Cys Ala Gln Gly Phe Gly Lys Ala Thr His  
                   65                  70                  75                  80  
 Gly Gly Lys Trp Gly Asp Ile Tyr Met Val Thr Ser Asp Gln Asp Asp  
                   85                  90                  95  
 40 Asp Val Val Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Thr Gln  
                   100                  105                  110  
 Asp Arg Pro Leu Trp Ile Ile Phe Gln Arg Asp Met Ile Ile Tyr Leu  
                   115                  120                  125  
 45 Gln Gln Glu Met Val Val Thr Ser Asp Thr Thr Ile Asp Gly Arg Gly  
                   130                  135                  140  
 Ala Lys Val Glu Leu Val Tyr Gly Gly Ile Thr Leu Met Asn Val Lys  
 50   145                  150                  155                  160  
 Asn Val Ile Ile His Asn Ile Asp Ile His Asp Val Arg Val Leu Pro  
                   165                  170                  175  
 55 Gly Gly Arg Ile Lys Ser Asn Gly Gly Pro Ala Ile Pro Arg His Gln  
                   180                  185                  190  
 Ser Asp Gly Asp Ala Ile His Val Thr Gly Ser Ser Asp Ile Trp Ile  
                   195                  200                  205

- 72 -

Asp His Cys Thr Leu Ser Lys Ser Phe Asp Gly Leu Val Asp Val Asn  
 210 215 220  
 5 Trp Gly Ser Thr Gly Val Thr Ile Ser Asn Cys Lys Phe Thr His His  
 225 230 235 240  
 Glu Lys Ala Val Leu Leu Gly Ala Ser Asp Thr His Phe Gln Asp Leu  
 245 250 255  
 10 Lys Met His Val Thr Leu Ala Tyr Asn Ile Phe Thr Asn Thr Val His  
 260 265 270  
 15 Glu Arg Met Pro Arg Cys Arg Phe Gly Phe Phe Gln Ile Val Asn Asn  
 275 280 285  
 Phe Tyr Asp Arg Trp Asp Lys Tyr Ala Ile Gly Gly Ser Ser Asn Pro  
 290 295 300  
 20 Thr Ile Leu Ser Gln Gly Asn Lys Phe Val Ala Pro Asp Phe Ile Tyr  
 305 310 315 320  
 Lys Lys Asn Val Cys Leu Arg Thr Gly Ala Gln Glu Pro Glu Trp Met  
 325 330 335  
 25 Thr Trp Asn Trp Arg Thr Gln Asn Asp Val Leu Glu Asn Gly Ala Ile  
 340 345 350  
 Phe Val Ala Ser Gly Ser Asp Pro Val Leu Thr Ala Glu Gln Asn Ala  
 355 360 365  
 30 Gly Met Met Gln Ala Glu Pro Gly Asp Met Val Pro Gln Leu Thr Met  
 370 375 380  
 35 Asn Ala Gly Val Leu Thr Cys Ser Pro Gly Ala Pro Cys  
 385 390 395

(2) INFORMATION FOR SEQ ID NO:11:

40

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 312 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: cDNA

50

- (ix) FEATURE:
- (A) NAME/KEY: CDS
  - (B) LOCATION: 1..276

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATG TCG ATT CTT GGA GGA ATT ACC GAA GTT AAA GAC AAT GAT AAC AGC  
 Met Ser Ile Leu Gly Gly Ile Thr Glu Val Lys Asp Asn Asp Asn Ser  
 1 5 10 15

- 73 -

GTC GAT TTC GAC GAG CTT GCT AAA TTC GCC ATC GCT GAA CAC AAC AAG 96  
 Val Asp Phe Asp Glu Leu Ala Lys Phe Ala Ile Ala Glu His Asn Lys  
 20 25 30  
 5 AAG GAG AAT GCT GCT CTG GAG TTT GGA AAA GTA ATA GAA AAA AAG CAG 144  
 Lys Glu Asn Ala Ala Leu Glu Phe Gly Lys Val Ile Glu Lys Lys Gln  
 35 40 45  
 10 CAG GCG GTA CAG GGC ACC ATG TAT TAT ATA AAA GTG GAA GCA AAT GAT 192  
 Gln Ala Val Gln Gly Thr Met Tyr Tyr Ile Lys Val Glu Ala Asn Asp  
 50 55 60  
 15 GGT GGT GAG AAG AAA ACT TAT GAA GCC AAG GTG TGG GTT AAG CTA TGG 240  
 Gly Gly Glu Lys Lys Thr Tyr Glu Ala Lys Val Trp Val Lys Leu Trp  
 65 70 75 80  
 GAA AAT TTC AAG GAA TTG CAG GAA CTC AAA CTT GTT TGATGTTGCC 286  
 Glu Asn Phe Lys Glu Leu Gln Glu Leu Lys Leu Val  
 20 85 90  
 ACCTCACCTTA ACTCCATATG GACGG 312

25 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 92 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

35 Met Ser Ile Leu Gly Gly Ile Thr Glu Val Lys Asp Asn Asp Asn Ser  
 1 5 10 15  
 40 Val Asp Phe Asp Glu Leu Ala Lys Phe Ala Ile Ala Glu His Asn Lys  
 20 25 30  
 Lys Glu Asn Ala Ala Leu Glu Phe Gly Lys Val Ile Glu Lys Lys Gln  
 35 40 45  
 45 Gln Ala Val Gln Gly Thr Met Tyr Tyr Ile Lys Val Glu Ala Asn Asp  
 50 55 60  
 Gly Gly Glu Lys Lys Thr Tyr Glu Ala Lys Val Trp Val Lys Leu Trp  
 65 70 75 80  
 50 Glu Asn Phe Lys Glu Leu Gln Glu Leu Lys Leu Val  
 85 90

55 (2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids

(B) TYPE: amino acid

- 74 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

10 Tyr Ile Leu Tyr Phe Thr Leu Ala Leu Val Thr Leu Leu Gln Pro Val  
 1 5 10 15

Arg Ser Ala Glu Asp Leu Gln Glu Ile Leu Pro  
 20 25  
 15

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:  
 20 (A) LENGTH: 28 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

30 Pro Val Arg Ser Ala Glu Asp Leu Gln Glu Ile Leu Pro Val Asn Glu  
 1 5 10 15

Thr Arg Arg Leu Thr Thr Ser Gly Ala Tyr Asn Ile  
 20 25  
 35

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:  
 40 (A) LENGTH: 24 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

50 Glu Thr Arg Arg Ser Leu Lys Thr Ser Gly Ala Tyr Asn Ile Ile Asp  
 1 5 10 15

Gly Cys Trp Arg Gly Lys Ala Asp  
 20  
 55

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 75 -

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Glu Thr Arg Arg Leu Thr Thr Ser Gly Ala Tyr Asn Ile Ile Asp Gly  
1 5 10 15  
Cys Trp Arg Gly Lys Ala Asp  
20

- 20 (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ala Tyr Asn Ile Ile Asp Gly Cys Trp Arg Gly Lys Ala Asp Trp Ala  
1 5 10 15  
Glu Asn Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly  
20 25

- 40 (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 45 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Arg Gly Lys Ala Asp Trp Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys  
1 5 10 15  
Ala Gln Gly

- 76 -

## (2) INFORMATION FOR SEQ ID NO:19:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- 10 (v) FRAGMENT TYPE: internal
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:
- 15 Gly Lys Ala Asp Trp Ala Glu Asn Arg Cys  
1 5 10

## 20 (2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
25 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
- Gly Lys Ala Asp Trp Ala Glu Asn Arg Lys  
35 1 5 10

## (2) INFORMATION FOR SEQ ID NO:21:

- 40 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
- Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly  
1 5 10

55

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids

- 77 -

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys Gly Thr Val Gly  
1 5 10 15

- Gly

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gly Phe Gly Lys Gly Thr Val Gly Gly Lys Asp Gly Asp Ile Tyr Ser  
1 5 10 15

Val Thr

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Gly Phe Gly Lys Gly Thr Val Gly Gly Lys Asp Gly Asp Ile Tyr Thr  
1 5 10 15

Val Thr

- 78 -

## (2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Lys Asp Gly Asp Ile Tyr Ser Val Thr Ser Glu Leu Asp Asp Asp Val  
1 5 10 15  
Ala

## (2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp Val  
1 5 10 15  
Ala

## (2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp  
1 5 10



- 79 -

## (2) INFORMATION FOR SEQ ID NO:28:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp Val Ala Asn  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:29:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp Val Ala Asn Pro Lys  
1 5 10 15

Glu

## (2) INFORMATION FOR SEQ ID NO:30:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ile Tyr Ser Val Thr Ser Glu Leu Asp Asp Asp Val Ala Asn Pro Lys  
1 5 10 15

Glu Gly Thr Leu  
20

- 80 -

## (2) INFORMATION FOR SEQ ID NO:31:

5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:  
Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp Val Ala Asn Pro Lys  
1 5 10 15  
Glu Gly Thr Leu  
20 20

## (2) INFORMATION FOR SEQ ID NO:32:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:  
Asn Pro Lys Glu Gly Thr Leu Arg Phe Gly Ala Ala Gln Asn Arg Pro  
1 5 10 15  
Leu Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu  
20 25

## (2) INFORMATION FOR SEQ ID NO:33:

45 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:  
Trp Ile Ile Phe Glu Arg Asp Met Val Ile Arg Leu Asp Lys Glu Met  
1 5 10 15

- 81 -

Val Val Asn Ser  
20

5

## (2) INFORMATION FOR SEQ ID NO:34:

## (i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

15

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

20

Leu Asp Lys Glu Met Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg  
1 5 10 15

Gly Ala Lys

25

## (2) INFORMATION FOR SEQ ID NO:35:

## (i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 26 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

35

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

40

Asp Lys Thr Ile Asp Gly Arg Gly Ala Lys Val Glu Ile Ile Asn Ala  
1 5 10 15

Gly Phe Thr Leu Asn Gly Val Lys Asn Val  
20 25

45

## (2) INFORMATION FOR SEQ ID NO:36:

50

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

55

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

- 82 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Gly Phe Thr Leu Asn Gly Val Lys Asn Val Ile Ile His Asn Ile Asn  
1 5 10 15  
Met His Asp Val Lys Val Asn Pro Gly Gly Leu  
20 25

10 (2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 amino acids  
(B) TYPE: amino acid  
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

His Asp Val Lys Val Asn Pro Gly Gly Leu Ile Lys Ser Asn Asp Gly  
1 5 10 15  
Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala  
20 25

30

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 amino acids  
35 (B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

Asn Pro Gly Gly Leu Ile Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg  
1 5 10 15  
Ala Gly Ser Asp Gly Asp Ala  
20

50

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
55 (B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 83 -

(v) FRAGMENT TYPE: internal

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Ile Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly  
1 5 10 15

10 Asp Ala

15 (2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala  
1 5 10

30

(2) INFORMATION FOR SEQ ID NO:41:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile  
1 5 10

50 (2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

- 84 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

5     Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser  
      1                    5                    10                    15  
  
      Ser Gln

10

(2) INFORMATION FOR SEQ ID NO:43:

15     (i) SEQUENCE CHARACTERISTICS:  
         (A) LENGTH: 23 amino acids  
         (B) TYPE: amino acid  
         (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

25     Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser  
      1                    5                    10                    15  
  
      Ser Gln Ile Trp Ile Asp His  
30                    20

(2) INFORMATION FOR SEQ ID NO:44:

35     (i) SEQUENCE CHARACTERISTICS:  
         (A) LENGTH: 28 amino acids  
         (B) TYPE: amino acid  
         (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

      Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser  
      1                    5                    10                    15  
  
50     Ser Gln Ile Trp Ile Asp His Cys Ser Leu Ser Lys  
                         20                    25

(2) INFORMATION FOR SEQ ID NO:45:

55

      (i) SEQUENCE CHARACTERISTICS:  
         (A) LENGTH: 28 amino acids  
         (B) TYPE: amino acid  
         (D) TOPOLOGY: linear

- 85 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Trp Ile Asp His Cys Ser Leu Ser Lys Ser Val Asp Gly Leu Val Asp  
1 5 10 15  
Ala Lys Leu Gly Thr Thr Arg Leu Thr Val Ser Asn  
20 25

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Leu Gly Thr Thr Arg Leu Thr Val Ser Asn Ser Leu Phe Thr Gln His  
1 5 10 15  
Gln Phe Val Leu Leu Phe Gly Ala Gly Asp Glu  
20 25

(2) INFORMATION FOR SEQ ID-NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Phe Val Leu Leu Phe Gly Ala Gly Asp Glu Asn Ile Glu Asp Arg Gly  
1 5 10 15  
Met Leu Ala Thr Val Ala Phe Asn Thr Phe Thr Asp  
20 25

(2) INFORMATION FOR SEQ ID NO:48:

- 86 -

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:
- Leu Ala Thr Val Ala Phe Asn Thr Phe Thr Asp Asn Val Asp Gln Arg  
1 5 10 15  
Met Pro
- 15
- (2) INFORMATION FOR SEQ ID NO:49:
- 20 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- 25 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:
- Phe Thr Asp Asn Val Asp Gln Arg Met Pro Arg Cys Arg His Gly Phe  
1 5 10 15  
Phe Gln Val Val  
20
- 35
- 40 (2) INFORMATION FOR SEQ ID NO:50:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:
- Arg Cys Arg His Gly Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys  
1 5 10 15  
Trp Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro  
20 25
- 55



- 87 -

## (2) INFORMATION FOR SEQ ID NO:51:

5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

His Gly Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys Trp Gly Ser  
1 5 10 15

20 Tyr Ala Ile Gly Gly Ser Ala Ser Pro  
20 25

## (2) INFORMATION FOR SEQ ID NO:52:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Phe Gln Val Val Asn Asn Asn Tyr Asp Lys Trp Gly Ser Tyr Ala Ile  
1 5 10 15

40 Gly Gly Ser Ala Ser Pro  
20

## (2) INFORMATION FOR SEQ ID NO:53:

45 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Val Asn Asn Asn Tyr Asp Lys Trp Gly Ser Tyr Ala Ile Gly Gly Ser  
1 5 10 15

## Ala Ser Pro

5

## (2) INFORMATION FOR SEQ ID NO:54:

## (i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

15

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

20

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr  
1 5 10

25

## (2) INFORMATION FOR SEQ ID NO:55:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

30

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15

40

## (2) INFORMATION FOR SEQ ID NO:56:

## (i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

50

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

55

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15

Gly Asn Arg Phe

20

## (2) INFORMATION FOR SEQ ID NO:57:

5

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15

20

Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys  
20 25

## 25 (2) INFORMATION FOR SEQ ID NO:58:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: amino acid

30

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys Lys Asn Val Leu Gly Arg  
1 5 10 15

40

His Gly Glu Ala Ala Ala Glu Ser Met Lys Trp Asn  
20 25

45

## (2) INFORMATION FOR SEQ ID NO:59:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

50

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

55

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

- 90 -

Glu Ala Ala Ala Glu Ser Met Lys Trp Asn Trp Arg Thr Asn Lys Asp  
 1 5 10 15

Val Leu Glu Asn Gly Ala Ile Phe Val Ala  
 5 20 25

## (2) INFORMATION FOR SEQ ID NO:60:

10 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 23 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide  
 (v) FRAGMENT TYPE: internal

## 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Glu Ser Met Lys Trp Asn Trp Arg Thr Asn Lys Asp Val Leu Glu Asn  
 1 5 10 15

25 Gly Ala Ile Phe Val Ala Ser  
 20

## (2) INFORMATION FOR SEQ ID NO:61:

30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide  
 (v) FRAGMENT TYPE: internal

## 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Glu Phe Gly Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile  
 1 5 10 15

45 Phe Val Ala Ser  
 20

## 50 (2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: peptide  
 (v) FRAGMENT TYPE: internal

- 91 -

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

5      Lys Trp Asn Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile  
      1                      5                      10                      15  
      Phe Val Ala Ser  
                            20

10

## (2) INFORMATION FOR SEQ ID NO:63:

15      - (i) SEQUENCE CHARACTERISTICS:  
            (A) LENGTH: 30 amino acids  
            (B) TYPE: amino acid  
            (D) TOPOLOGY: linear

20      (ii) MOLECULE TYPE: peptide

      (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

25      Lys Trp Asn Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile  
      1                      5                      10                      15  
      Phe Val Ala Ser Gly Val Asp Pro Val Leu Thr Pro Glu Gln  
30                      20                      25                      30

## (2) INFORMATION FOR SEQ ID NO:64:

35      (i) SEQUENCE CHARACTERISTICS:  
            (A) LENGTH: 17 amino acids  
            (B) TYPE: amino acid  
            (D) TOPOLOGY: linear

40      (ii) MOLECULE TYPE: peptide

      (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

45      Trp Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile Phe Val Ala  
      1                      5                      10                      15  
50      Ser

## (2) INFORMATION FOR SEQ ID NO:65:

55      (i) SEQUENCE CHARACTERISTICS:  
            (A) LENGTH: 14 amino acids  
            (B) TYPE: amino acid  
            (D) TOPOLOGY: linear

- 92 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Asn Lys Asp Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser  
1 5 10

10

(2) INFORMATION FOR SEQ ID NO:66:

15

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser  
1 5 10

30

(2) INFORMATION FOR SEQ ID NO:67:

35

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

45

Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly  
1 5 10

(2) INFORMATION FOR SEQ ID NO:68:

50

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 14 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro  
5           1                 5                                 10

(2) INFORMATION FOR SEQ ID NO:69:

10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val Leu  
1                    5                         10                         15

25 Thr

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val Leu  
1 5 10 15

Thr Pro Glu Gln  
20

50 (2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

5 Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro  
1 5 10 15  
Ala Glu Pro Gly  
20

10

(2) INFORMATION FOR SEQ ID NO:72:

15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Gly Val Asp Pro Val Leu Thr Pro Glu  
1 5

30 (2) INFORMATION FOR SEQ ID NO:73:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

45 Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala  
1 5 10

(2) INFORMATION FOR SEQ ID NO:74:

50 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 16 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal



- 95 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

5           Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro  
          1                   5                   10                   15

(2) INFORMATION FOR SEQ ID NO:75:

10       (i) SEQUENCE CHARACTERISTICS:  
          (A) LENGTH: 20 amino acids  
          (B) TYPE: amino acid  
          (D) TOPOLOGY: linear

15       (ii) MOLECULE TYPE: peptide

          (v) FRAGMENT TYPE: internal

20       (xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

          Ser Ala Gly Met Ile Pro Ala Glu Pro Gly Glu Ser Ala Leu Ser Leu  
          1                   5                   10                   15  
  
25       Thr Ser Ser Ala  
                          20

(2) INFORMATION FOR SEQ ID NO:76:

30       (i) SEQUENCE CHARACTERISTICS:  
          (A) LENGTH: 18 amino acids  
          (B) TYPE: amino acid  
          (D) TOPOLOGY: linear

35       (ii) MOLECULE TYPE: peptide

          (v) FRAGMENT TYPE: internal

40       (xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

          Glu Pro Gly Glu Ser Ala Leu Ser Leu Thr Ser Ser Ala Gly Val Leu  
          1                   5                   10                   15  
  
45       Ser Cys

(2) INFORMATION FOR SEQ ID NO:77:

50       (i) SEQUENCE CHARACTERISTICS:  
          (A) LENGTH: 15 amino acids  
          (B) TYPE: amino acid  
          (D) TOPOLOGY: linear

55       (ii) MOLECULE TYPE: peptide

          (v) FRAGMENT TYPE: internal

- 96 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Thr Ser Ser Ala Gly Val Leu Ser Cys Gln Pro Gly Ala Pro Cys  
1 5 10 15

5

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

15

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp His Val Asp Gln Arg  
1 5 10 15

20

Met Pro

25

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

35

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

Tyr Ile Leu Tyr Phe Thr Leu Ala Leu Val Ala Leu Leu Gln Pro Val  
1 5 10 15

40

Arg Ser Ala Glu Gly Val Gly Glu Ile Leu Pro  
20 25

45

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

55

(v) FRAGMENT TYPE: internal

- 97 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

5       Leu Asn Gln Glu Leu Val Val Asn Ser Asp Lys Thr Ile Asp Gly Arg  
      1                       5                       10                       15  
      Gly Val Lys

10

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

15       -       (A) LENGTH: 21 amino acids  
              (B) TYPE: amino acid  
              (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

25       Gly Met Leu Ala Thr Val Ala Phe Asn Met Phe Thr Asp Asn Val Asp  
      1                       5                       10                       15  
      Gln Arg Met Pro Arg  
                              20

30

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

35       (A) LENGTH: 28 amino acids  
              (B) TYPE: amino acid  
              (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

45

Arg Gly Lys Ala Asp Trp Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys  
1                       5                       10                       15

50

Ala Gln Gly Phe Gly Lys Gly Thr Val Gly Gly Lys  
                              20                       25

(2) INFORMATION FOR SEQ ID NO:83:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

- 98 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys  
1 5 10 15

10

Gly Thr Val Gly Gly Lys  
20

15 (2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 amino acids

(B) TYPE: amino acid

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys  
1 5 10 15

30

Gly Thr Val Gly Gly Lys Asp Gly Asp  
20 25

35

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids

40

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

45

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

50

Gly Phe Gly Lys Gly Thr Val Gly Gly Lys Asp Gly Asp Ile Tyr Thr  
1 5 10 15

Val Thr Ser Glu Leu Asp Asp Asp Val Ala Asn  
20 25

55

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:

- 99 -

(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

Lys Gly Thr Val Gly Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser  
-1 5 10 15

15

Glu Leu Asp Asp Asp Val Ala Asn  
20

(2) INFORMATION FOR SEQ ID NO:87:

20

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

35

Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu Asp Asp Asp  
1 5 10 15

Val Ala Asn

40 (2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

55

Glu Asn Asn Arg Gln Ala Leu Ala Asp Cys Ala Gln Gly Phe Ala Lys  
1 5 10 15

Gly Thr Tyr Gly Gly Lys Trp Gly Asp  
20 25

- 100 -

## (2) INFORMATION FOR SEQ ID NO:89:

5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

## 15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Lys Gly Thr Tyr Gly Gly Lys Trp Gly Asp Val Tyr Thr Val Thr Ser  
1 5 10 15

20 Asn Leu Asp Asp Asp Val Ala Asn  
20

## (2) INFORMATION FOR SEQ ID NO:90:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

## 35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp  
1 5 10 15

40 Ala Ile Ser Ile Ser Gly  
20

## (2) INFORMATION FOR SEQ ID NO:91:

45 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 25 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

55

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp

- 101 -

1 5 10 15

Ala Ile Ser Ile Ser Gly Ser Ser Gln  
20 25

5

## (2) INFORMATION FOR SEQ ID NO:92:

## (i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 30 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

15

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

20

Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp  
1 5 10 15

25

Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His  
20 25 30

## (2) INFORMATION FOR SEQ ID NO:93:

## (i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 35 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

35

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

40

Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp  
1 5 10 15

45

Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Cys Ser  
20 25 30

Leu Ser Lys  
35

## 50 (2) INFORMATION FOR SEQ ID NO:94:

## (i) SEQUENCE CHARACTERISTICS:

55

- (A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

- 102 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

5 Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile  
1 5 10 15  
Ser Gly

10

(2) INFORMATION FOR SEQ ID NO:95:

15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20

(v) FRAGMENT TYPE: internal

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile  
1 5 10 15

30

Ser Gly Ser Ser Gln  
20

(2) INFORMATION FOR SEQ ID NO:96:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

Lys Lys Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
1 5 10 15

50

Gly Ser Ser Gln Ile Trp Ile Asp His Lys Lys  
20 25

(2) INFORMATION FOR SEQ ID NO:97:

55

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 31 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear



- 103 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

10 Lys Lys Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
1 5 10 15  
Gly Ser Ser Gln Ile Trp Ile Asp His Cys Ser Leu Ser Lys Lys  
20 25 30

15

(2) INFORMATION FOR SEQ ID NO:98:

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 31 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

25

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

30

Lys Lys Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
1 5 10 15  
Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser Lys Lys  
20 25 30

35

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:

40

(A) LENGTH: 31 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

45

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

50

Lys Lys Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
1 5 10 15  
Gly Ser Ser Gln Ile Trp Ile Asp His Leu Ser Leu Ser Lys Lys  
20 25 30

55

(2) INFORMATION FOR SEQ ID NO:100:

- 104 -

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 31 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

15 Lys Lys Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
1 5 10 15  
Gly Ser Ser Gln Ile Trp Ile Asp His Glu Ser Leu Ser Lys Lys  
20 25 30

20 (2) INFORMATION FOR SEQ ID NO:101:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

35 Gly Pro Pro Ile Leu Arg Gln Ala Ser Asp Gly Asp Thr Ile Asn Val  
1 5 10 15  
Ala Gly Ser Ser Gln  
20

40

(2) INFORMATION FOR SEQ ID NO:102:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: peptide

50

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

55

Arg His Gly Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys Trp Gly  
1 5 10 15  
Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr

- 105 -

20

25

## (2) INFORMATION FOR SEQ ID NO:103:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys Trp Gly Ser Tyr Ala  
1 5 10 15  
Ile Gly Gly Ser Ala Ser Pro Thr  
20

25 (2) INFORMATION FOR SEQ ID NO:104:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15  
Gly Asn Arg Phe Cys Ala Pro Asp  
20

45 (2) INFORMATION FOR SEQ ID NO:105:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

- 106 -

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15

5 Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg  
20 25

(2) INFORMATION FOR SEQ ID NO:106:

-10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln  
1 5 10 15  
25 Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys  
20 25

(2) INFORMATION FOR SEQ ID NO:107:

30 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Arg Trp Gly Thr Tyr Ala  
1 5 10 15  
45 Ile Gly Gly Ser Ser Ala Pro Thr  
20

(2) INFORMATION FOR SEQ ID NO:108:

50 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: peptide  
(v) FRAGMENT TYPE: internal

- 107 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

5 Gly Thr Tyr Ala Ile Gly Gly Ser Ser Ala Pro Thr Ile Leu Cys Gln  
1 5 10 15  
Gly Asn Arg Phe Leu Ala Pro Asp Asp Gln Ile Lys  
20 25

10

(2) INFORMATION FOR SEQ ID NO:109:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 29 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

25

Lys Asp Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp  
1 5 10 15

30

Pro Val Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro  
20 25

(2) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 26 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

45

Lys Asp Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp  
1 5 10 15

50

Pro Val Leu Thr Pro Glu Gln Ser Ala Gly  
20 25

(2) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

55

(A) LENGTH: 31 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

- 108 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
1 5 10 15  
Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro Ala Glu Pro Gly  
20 25 30

15 (2) INFORMATION FOR SEQ ID NO:112:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
1 5 10 15  
Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro Ala Glu  
20 25

35

(2) INFORMATION FOR SEQ ID NO:113:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
1 5 10 15  
Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro  
20 25

55

(2) INFORMATION FOR SEQ ID NO:114:

(i) SEQUENCE CHARACTERISTICS:

- 109 -

- (A) LENGTH: 24 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
—1                    5                    10                    15  
Leu Thr Pro Glu Gln Ser Ala Gly  
                    20

- 20 (2) INFORMATION FOR SEQ ID NO:115:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
35 1                    5                    10                    15  
Leu Thr Pro Glu Gln  
                    20

- 40 (2) INFORMATION FOR SEQ ID NO:116:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 45 (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal

- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

Leu Leu Glu Asn Gly Ala Ile Phe Val Thr Ser Gly Ser Asp Pro Val  
55 1                    5                    10                    15  
Leu Thr Pro Val Gln Ser Ala Gly Met Ile Pro  
                    20                    25

- 110 -

## (2) INFORMATION FOR SEQ ID NO:117:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 60 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

Ala Tyr Asn Ile Ile Asp Gly Cys Trp Arg Gly Lys Ala Asp Trp Ala  
1 5 10 15  
Glu Asn Arg Lys Ala Leu Ala Asp Cys Ala Gln Gly Phe Gly Lys Gly  
20 25 30  
Thr Val Gly Gly Lys Asp Gly Asp Ile Tyr Thr Val Thr Ser Glu Leu  
35 40 45  
25 Asp Asp Asp Val Ala Asn Pro Lys Glu Gly Thr Leu  
50 55 60

30 (2) INFORMATION FOR SEQ ID NO:118:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 46 amino acids  
(B) TYPE: amino acid  
35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

45 His Asp Val Lys Val Asn Pro Gly Gly Leu Ile Lys Ser Asn Asp Gly  
1 5 10 15  
Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp Ala Ile Ser Ile Ser  
20 25 30  
50 Gly Ser Ser Gln Ile Trp Ile Asp His Cys Ser Leu Ser Lys  
35 40 45

55 (2) INFORMATION FOR SEQ ID NO:119:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 amino acids  
(B) TYPE: amino acid



- 111 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

10

Arg Cys Arg His Gly Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys  
1 5 10 15

Trp Gly Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser  
15 20 25 30

Gln Gly Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys  
35 40 45

20

(2) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 47 amino acids  
25 (B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

30 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

35 Glu Ala Ala Ala Glu Ser Met Lys Trp Asn Trp Arg Thr Asn Lys Asp  
1 5 10 15

Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp Pro Val  
20 25 30

40

Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro Ala Glu Pro Gly  
35 40 45

45 (2) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 45 amino acids  
50 (B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

55 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

Arg Gly Lys Ala Asp Trp Ala Glu Asn Arg Lys Ala Leu Ala Asp Cys

- 112 -

10 (2) INFORMATION FOR SEQ ID NO:122:

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

25 Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp  
1 5 10 15

Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Cys Ser  
20 25 30

30 Leu Ser Lys  
35

(2) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

Arg His Gly Phe Phe Gln Val Val Asn Asn Asn Tyr Asp Lys Trp Gly  
1 5 10 15  
50 Ser Tyr Ala Ile Gly Gly Ser Ala Ser Pro Thr Ile Leu Ser Gln Gly  
20 25 30  
Asn Arg Phe Cys Ala Pro Asp Glu Arg Ser Lys  
55 35 40

(2) INFORMATION FOR SEQ ID NO:124:

- 113 -

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

10

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

Arg Thr Asn Lys Asp Val Leu Glu Asn Gly Ala Ile Phe Val Ala Ser  
1\_ 5 10 15

15

Gly Val Asp Pro Val Leu Thr Pro Glu Gln Ser Ala Gly Met Ile Pro  
20 25 30

20

Ala Glu Pro Gly  
35

## (2) INFORMATION FOR SEQ ID NO:125:

25

## (i) SEQUENCE CHARACTERS:

- (A) LENGHT 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30

(ii) MOLECULAR TYPE: peptide

(v) FRAGMENT TYPE: internal

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

Ser Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile  
1 5 10 15

40

Asp His Cys Ser Leu Ser Lys Ser  
20

## (2) INFORMATION FOR SEQ ID NO:126:

45

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser Lys Ser

- 114 -

1

5

10

15

## (2) INFORMATION FOR SEQ ID NO:127:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser  
1 5 10 15

20

Lys Ser

## (2) INFORMATION FOR SEQ ID NO:128:

25

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

Asp Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser  
1 5 10 15

40

Ser Leu Ser Lys Ser  
20

## (2) INFORMATION FOR SEQ ID NO:129:

45

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

- 115 -

Ser Asp Gly Asp Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile  
1 5 10 15

Asp His Ser Ser Leu Ser Lys Ser  
20

(2) INFORMATION FOR SEQ ID NO:130:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

\_(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:

Ser Glu Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser  
1 5 10 15

Lys Ser

(2) INFORMATION FOR SEQ ID NO:131:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

Ser Lys Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser  
1 5 10 15

Lys Ser

(2) INFORMATION FOR SEQ ID NO:132:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

- 116 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

5 Asp Lys Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser  
1 5 10 15  
Lys Glu

10

(2) INFORMATION FOR SEQ ID NO:133:

15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

Asp Lys Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser  
1 5 10 15  
Leu Ser Lys Glu  
20

30

(2) INFORMATION FOR SEQ ID NO:134:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

Asp Lys Glu Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu  
1 5 10 15  
Ser Lys Glu

50

(2) INFORMATION FOR SEQ ID NO:135:

55

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 18 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

- 117 -

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

10 Asp Lys Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser  
1 5 10 15  
Lys Lys

15

(2) INFORMATION FOR SEQ ID NO:136:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

20

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

25

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:136:

30

Asp Lys Glu Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu  
1 5 10 15

Ser Lys Glu Lys  
20

35

(2) INFORMATION FOR SEQ ID NO:137:

(i) SEQUENCE CHARACTERISTICS:

40

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

45

(v) FRAGMENT TYPE: internal

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

Ser Gly Ser Ser Gln Ile Ala Ile Asp His Ser Ser Leu Ser Lys Ser  
1 5 10 15

55

(2) INFORMATION FOR SEQ ID NO:138:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

- 118 -

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5

(v) FRAGMENT TYPE: internal

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

Asp Lys Glu Ser Gly Ser Ser Gln Ile Ala Ile Asp His Ser Ser Leu  
1 5 10 15

15

Ser Lys Ser Glu  
20

(2) INFORMATION FOR SEQ ID NO:139:

20

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

Asp Lys Glu Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser Leu  
1 5 10 15

35

Ser Lys Glu Lys Asp  
20

(2) INFORMATION FOR SEQ ID NO:140:

40

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

Ser Gln Ile Trp Ile Asp His Ser Ser Leu Ser Lys Ser  
1 5 10

55

(2) INFORMATION FOR SEQ ID NO:141:

(i) SEQUENCE CHARACTERISTICS:



- 119 -

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

Asp Lys Glu Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser  
1 5 10 15  
Ser Leu Ser Lys Glu Lys Asp  
20

15

(2) INFORMATION FOR SEQ ID NO:142:

20

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Glu  
1 5 10 15  
Pro Val Leu Thr Pro His Gln Ser Ala Gly  
20 25

35

(2) INFORMATION FOR SEQ ID NO:143:

40

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asn  
1 5 10 15  
Pro Val Leu Thr Pro His Gln Ser Ala Gly  
20 25

55

(2) INFORMATION FOR SEQ ID NO:144:

- 120 -

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGHT: 26 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULAR TYPE: peptide
- (v) FRAGMENT TYPE: internal
- (xi) SEQUENCE DESCRIPTION SEQ ID NO:144:
- Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Ala  
1 5 10 15  
Pro Val Leu Thr Pro His Gln Ser Ala Gly  
20 25
- (2) INFORMATION FOR SEQ ID NO:145:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:
- Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Ser  
1 5 10 15  
Pro Val Leu Thr Pro His Gln Ser Ala Gly  
20 25
- (2) INFORMATION FOR SEQ ID NO:146:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (v) FRAGMENT TYPE: internal
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:
- Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Gln  
1 5 10 15

- 121 -

Pro Val Leu Thr Pro His Gln Ser Ala Gly  
20 25

## (2) INFORMATION FOR SEQ ID NO:147:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:147:

Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp  
1 5 10 15

Ala Val Leu Thr Pro His Gln Ser Ala Gly  
20 25

## (2) INFORMATION FOR SEQ ID NO:148:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:148:

Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp  
1 5 10 15

Ser Val Leu Thr Pro His Gln Ser Ala Gly  
20 25

## (2) INFORMATION FOR SEQ ID NO:149:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:149:

- 122 -

Lys Asp Val Leu His Asn Gly Ala Ile Phe Val Ala Ser Gly Val Asp  
 1 5 10 15

5 Gly Val Leu Thr Pro His Gln Ser Ala Gly  
 20 25

## (2) INFORMATION FOR SEQ ID NO:150:

10 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 35 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide  
 (v) FRAGMENT TYPE: internal

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:150:

Lys Ser Asn Asp Gly Pro Ala Ala Pro Arg Ala Gly Ser Asp Gly Asp  
 1 5 10 15

25 Ala Ile Ser Ile Ser Gly Ser Ser Gln Ile Trp Ile Asp His Ser Ser  
 20 25 30

30 Leu Ser Lys  
 35

**Claims**

1. All or a portion of an isolated peptide of *Amb a* I.1, said peptide or portion thereof comprising at least one T cell epitope of *Amb a* I.1, said peptide comprising an amino acid sequence selected from the group consisting of:
- a) AMB 1-1.1 (SEQ ID NO:85);
  - b) AMB 1-2.1 (SEQ ID NO:86);
  - c) AMB 1-3.1 (SEQ ID NO:87);
  - d) AMB 1-4.1 (SEQ ID NO:84);
  - e) AMB 1-5.1 (SEQ ID NO:83);
  - f) AMB 1-6.1 (SEQ ID NO:82);
  - g) AMB 2-4.1 (SEQ ID NO:90);
  - h) AMB 2-3.1 (SEQ ID NO:91);
  - i) AMB 2-5.1 (SEQ ID NO:92);
  - j) AMB 2-6.1 (SEQ ID NO:93);
  - k) AMB 2-2.1 (SEQ ID NO:94);
  - l) AMB 2-1.1 (SEQ ID NO:95);
  - m) RAE 70.1-1 (SEQ ID NO:43);
  - n) AMB 2-7.1 (SEQ ID NO:96);
  - o) AMB 2-8.1 (SEQ ID NO:97);
  - p) AMB 2-9.1 (SEQ ID NO:98);
  - q) AMB 2-10.1 (SEQ ID NO:99);
  - r) AMB 2-11.1 (SEQ ID NO:100);
  - s) RAE 70.1 (SEQ ID NO:44);
  - t) AMB 3-4.1 (SEQ ID NO:103);
  - u) AMB 3-5.1 (SEQ ID NO:102);
  - v) AMB 3-3.1 (SEQ ID NO:104);
  - w) AMB 3-2.1 (SEQ ID NO:105);
  - x) AMB 3-1.1 (SEQ ID NO:106);
  - y) AMB 4-8.1 (SEQ ID NO:109);
  - z) AMB 4-9.1 (SEQ ID NO:110);
  - a') AMB 4-6.1 (SEQ ID NO:111);
  - b') AMB 4-5.1 (SEQ ID NO:112);
  - c') AMB 4-3.1 (SEQ ID NO:113);
  - d') AMB 4-2.1 (SEQ ID NO:114);
  - e') AMB 4-1.1 (SEQ ID NO:115);
  - f) AMB 2-38.1 (SEQ ID NO:141);
  - g') RA-02.1 (SEQ ID NO:150); and
  - h') AMB 2-36.1 (SEQ ID NO:139).

2. All or a portion of an isolated peptide of claim 1 which has a stimulation index of at least 2.0.

5 3. All or a portion of an isolated peptide of claim 1 which has minimal immunoglobulin E stimulating activity.

10 4. All or a portion of an isolated peptide of claim 1 which does not bind immunoglobulin E specific for *Amb a* I.1 in a substantial percentage of individuals sensitive to *Amb a* I.1, or if binding of the peptide to said immunoglobulin E occurs, such binding does not result in release of mediators from mast cells or basophils in a substantial percentage of individuals sensitive to *Amb a* I.1.

15 5. All or a portion of an isolated peptide of claim 1 which binds immunoglobulin E to a substantially lesser extent than *Amb a* I.1 binds said immunoglobulin E.

20 6. All or a portion of an isolated peptide of claim 1 which modifies, in a ragweed pollen-sensitive individual to whom it is administered, the allergic response of the individual to ragweed pollen.

7. A portion of an isolated peptide of claim 1 wherein the portion comprises at least fifteen amino acid residues.

25 8. An isolated peptide of *Amb a* I.1 comprising an amino acid sequence selected from the group consisting of:

- a) AMB 1-1.1 (SEQ ID NO:85);
- b) AMB 1-2.1 (SEQ ID NO:86);
- c) AMB 1-3.1 (SEQ ID NO:87);
- d) AMB 1-4.1 (SEQ ID NO:84);
- 30 e) AMB 1-5.1 (SEQ ID NO:83);
- f) AMB 1-6.1 (SEQ ID NO:82);
- g) AMB 2-4.1 (SEQ ID NO:90);
- h) AMB 2-3.1 (SEQ ID NO:91);
- i) AMB 2-5.1 (SEQ ID NO:92);
- 35 j) AMB 2-6.1 (SEQ ID NO:93);
- k) AMB 2-2.1 (SEQ ID NO:94);
- l) AMB 2-1.1 (SEQ ID NO:95);
- m) RAE 70.1-1 (SEQ ID NO:43);
- n) AMB 2-7.1 (SEQ ID NO:96);

- 5 o) AMB 2-8.1 (SEQ ID NO:97);  
p) AMB 2-9.1 (SEQ ID NO:98);  
q) AMB 2-10.1 (SEQ ID NO:99);  
r) AMB 2-11.1 (SEQ ID NO:100);  
s) RAE 70.1 (SEQ ID NO:44);  
t) AMB 3-4.1 (SEQ ID NO:103);  
u) AMB 3-5.1 (SEQ ID NO:102);  
v) AMB 3-3.1 (SEQ ID NO:104);  
w) AMB 3-2.1 (SEQ ID NO:105);  
10 x) AMB 3-1.1 (SEQ ID NO:106);  
y) AMB 4-8.1 (SEQ ID NO:109);  
z) AMB 4-9.1 (SEQ ID NO:110);  
a') AMB 4-6.1 (SEQ ID NO:111);  
b') AMB 4-5.1 (SEQ ID NO:112);  
15 c') AMB 4-3.1 (SEQ ID NO:113);  
d') AMB 4-2.1 (SEQ ID NO:114);  
e') AMB 4-1.1 (SEQ ID NO:115);  
f) AMB 2-38.1 (SEQ ID NO:141);  
g') RA-02.1 (SEQ ID NO:150); and  
20 h') AMB 2-36.1 (SEQ ID NO:139).

9. All or a portion of an isolated peptide of claim 1, wherein said peptide or a portion thereof comprises at least two T cell epitopes *Amb a* 1.1.

25 10. An isolated nucleic acid having a sequence encoding all or a portion of a peptide of claim 1 or the functional equivalent of said nucleic acid sequence.

11. An isolated peptide which is immunologically cross-reactive with antibodies specific for all or a portion of a peptide of claim 1.

30

12. An isolated peptide which is immunologically cross-reactive with T cells reactive with all or a portion of a peptide of claim 1.

13. All or a portion of a modified peptide of claim 1.

35

14. A modified peptide of claim 8.

15. All or a portion of a modified peptide of claim 13 which has minimal immunoglobulin E stimulating activity.

16. All or a portion of a modified peptide of claim 13 which does not bind immunoglobulin E specific for *Amb a* I.1 in a substantial percentage of individuals sensitive to *Amb a* I.1, or if binding of the peptide to said immunoglobulin E occurs, such binding does not result in release of mediators from mast cells or basophils in a substantial percentage of individuals sensitive to *Amb a* I.1.

17. All or a portion of a modified peptide of claim 13 which binds immunoglobulin E to a substantially lesser extent than *Amb a* I.1 binds said immunoglobulin E.

18. All or a portion of a modified peptide of claim 13 which modifies, in a ragweed pollen-sensitive individual to whom it is administered, the allergic response of the individual to *Amb a* I.1.

15

19. All or a portion of an isolated peptide of *Ambrosia artemisiifolia*, said peptide or portion thereof comprising at least one T cell epitope of *Ambrosia artemisiifolia*, said peptide comprising an amino acid sequence selected from the group consisting of:

- a) RAE 67.1 (SEQ ID NO:13);
- 20 b) RAE 57.1 (SEQ ID NO:14);
- c) RAE 24.E (SEQ ID NO:15);
- d) RAE 24.1 (SEQ ID NO:16);
- e) RAE 22.E (SEQ ID NO:17);
- f) RAE 22.E-1 (SEQ ID NO:18);
- 25 g) RAE 3.1 (SEQ ID NO:20);
- h) RAE 22.E-2 (SEQ ID NO:21);
- i) RAE 5.D (SEQ ID NO:22);
- j) RAE 6.D (SEQ ID NO:23);
- k) RAE 6.1 (SEQ ID NO:24);
- 30 l) RAE 7.D (SEQ ID NO:25);
- m) RAE 7.D-1 (SEQ ID NO:26);
- n) RAE 40.1-6 (SEQ ID NO:27);
- o) RAE 40.1-5 (SEQ ID NO:28);
- p) RAE 40.1-4 (SEQ ID NO:29);
- 35 q) RAE 40.D (SEQ ID NO:30);
- r) RAE 40.1 (SEQ ID NO:31);
- s) RAE 61.1 (SEQ ID NO:32);
- t) RAE 80.1 (SEQ ID NO:33);
- u) RAE 45.1 (SEQ ID NO:34);



- v) RAE 75.1 (SEQ ID NO:35);  
w) RAE 62.1 (SEQ ID NO:36);  
x) RAE 69.1 (SEQ ID NO:37);  
y) RAE 69.1-1 (SEQ ID NO:38);  
5 z) RAE 69.1-2 (SEQ ID NO:39);  
a') RAE 69.1-3 (SEQ ID NO:40);  
b') RAE 70.1-3 (SEQ ID NO:41);  
c') RAE 70.1-2 (SEQ ID NO:42);  
d') RAE 71.1 (SEQ ID NO:45);  
10 e') RAE 65.1 (SEQ ID NO:46);  
f) RAE 63.1 (SEQ ID NO:47);  
g') RAE 76.1 (SEQ ID NO:48);  
h') RAE 27.1 (SEQ ID NO:49);  
i') RAE 66.1 (SEQ ID NO:50);  
15 j') RAE 66.1-1 (SEQ ID NO:51);  
k') RAE 66.1-2 (SEQ ID NO:52);  
l') RAE 66.1-3 (SEQ ID NO:53);  
m') RAE 64.1-3 (SEQ ID NO:54);  
n') RAE 64.1-2 (SEQ ID NO:55);  
20 o') RAE 64.1-1 (SEQ ID NO:56);  
p') RAE 64.1 (SEQ ID NO:57);  
q') RAE 73.1 (SEQ ID NO:58);  
r') RAE 74.1 (SEQ ID NO:59);  
s') RAE 74.1-1 (SEQ ID NO:60);  
25 t') RAE 29.1 (SEQ ID NO:61);  
u') RAE 29.1-1 (SEQ ID NO:62);  
v') RAE 28+29 (SEQ ID NO:63);  
w') RAE 29.1-2 (SEQ ID NO:64);  
x') RAE 29.1-3 (SEQ ID NO:65);  
30 y') RAE 29.1-4 (SEQ ID NO:66);  
z') RAE 28.1-3 (SEQ ID NO:67);  
a'') RAE 28.1-2 (SEQ ID NO:68);  
b'') RAE 28.1-1 (SEQ ID NO:69);  
c'') RAE 28.1 (SEQ ID NO:70);  
35 d'') RAE 20.1 (SEQ ID NO:71);  
e'') RAE 20.1-3 (SEQ ID NO:72);  
f'') RAE 20.1-2 (SEQ ID NO:73);  
g'') RAE 20.1-1 (SEQ ID NO:74);  
h'') RAE 21.1 (SEQ ID NO:75);

- 5 i") RAE 17.1 (SEQ ID NO:76);  
j") RAE 55.1 (SEQ ID NO:77);  
k") RAE 76.6 (SEQ ID NO:78);  
l") RAE 67.15 (SEQ ID NO:79);  
m") RAE 45.15 (SEQ ID NO:80); and  
n") RAE 27.15 (SEQ ID NO:81).

10 20. An isolated peptide comprising at least two regions, each region comprising at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia*, said regions derived from the same or from different protein allergens of *Ambrosia artemisiifolia*, said regions each comprising all or a portion of an amino acid sequence selected from the group consisting of:

- 15 a) AMB 1-1.1 (SEQ ID NO:85);  
b) AMB 1-2.1 (SEQ ID NO:86);  
c) AMB 1-3.1 (SEQ ID NO:87);  
d) AMB 1-4.1 (SEQ ID NO:84);  
e) AMB 1-5.1 (SEQ ID NO:83);  
f) AMB 1-6.1 (SEQ ID NO:82);  
20 g) AMB 2-4.1 (SEQ ID NO:90);  
h) AMB 2-3.1 (SEQ ID NO:91);  
i) AMB 2-5.1 (SEQ ID NO:92);  
j) AMB 2-6.1 (SEQ ID NO:93);  
k) AMB 2-2.1 (SEQ ID NO:94);  
25 l) AMB 2-1.1 (SEQ ID NO:95);  
m) RAE 70.1-1 (SEQ ID NO:43);  
n) AMB 2-7.1 (SEQ ID NO:96);  
o) AMB 2-8.1 (SEQ ID NO:97);  
p) AMB 2-9.1 (SEQ ID NO:98);  
30 q) AMB 2-10.1 (SEQ ID NO:99);  
r) AMB 2-11.1 (SEQ ID NO:100);  
s) RAE 70.1 (SEQ ID NO:44);  
t) AMB 3-4.1 (SEQ ID NO:103);  
u) AMB 3-5.1 (SEQ ID NO:102);  
35 v) AMB 3-3.1 (SEQ ID NO:104);  
w) AMB 3-2.1 (SEQ ID NO:105);  
x) AMB 3-1.1 (SEQ ID NO:106);  
y) AMB 4-8.1 (SEQ ID NO:109);  
z) AMB 4-9.1 (SEQ ID NO:110);

- 5
- a') AMB 4-6.1 (SEQ ID NO:111);
  - b') AMB 4-5.1 (SEQ ID NO:112);
  - c') AMB 4-3.1 (SEQ ID NO:113);
  - d') AMB 4-2.1 (SEQ ID NO:114);
  - e') AMB 4-1.1 (SEQ ID NO:115);
  - f') AMB 2-38.1 (SEQ ID NO:141);
  - g') RA-02.1 (SEQ ID NO:150); and
  - h') AMB 2-36.1 (SEQ ID NO:139).

- 10        21.    An isolated peptide of claim 20 wherein said peptide comprises various combinations of regions, said combinations of regions selected from the group consisting of:
- 15
- a) AMB 4-6.1 and RAE 70.1 (SEQ ID NO:111 and SEQ ID NO:44);
  - b) AMB 4-6.1 and AMB 2-5.1 (SEQ ID NO:111 and SEQ ID NO:92);
  - c) AMB 4-9.1 and AMB 2-5.1 (SEQ ID NO:110 and SEQ ID NO:92);
  - d) AMB 4-9.1 and RAE 70.1 (SEQ. ID NO:110 and SEQ ID NO:44);
  - e) AMB 4-6.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:92 and SEQ ID NO:86);
  - 20        f) AMB 4-9.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:92 and SEQ ID NO:86);
  - g) AMB 4-6.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:44 and SEQ ID NO:86);
  - h) AMB 4-9.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:44 and SEQ ID NO:86);
  - 25        i) AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102);
  - j) AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102);
  - 30        k) AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102);
  - l) AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102);
  - 35        m) AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103);
  - n) AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103);

- 5
- o) AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103);
- p) AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103);
- 10 q') AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:101, and SEQ ID NO:116);
- r') AMB 1-2.15, AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:89, SEQ ID NO:101, and SEQ ID NO:116);
- and
- s') AMB 1-2.15, AMB 2-1.15, AMB 4-3.15 and AMB 3-4.15 (SEQ ID NO:89, SEQ ID NO:101, SEQ ID NO:116 and SEQ ID NO:107).

15

22. An isolated peptide of claim 20, wherein the regions each comprise an amino acid sequence selected from the group consisting of:

- 20
- a) AMB 1-1.1 (SEQ ID NO:85);
- b) AMB 1-2.1 (SEQ ID NO:86);
- c) AMB 1-3.1 (SEQ ID NO:87);
- d) AMB 1-4.1 (SEQ ID NO:84);
- e) AMB 1-5.1 (SEQ ID NO:83);
- f) AMB 1-6.1 (SEQ ID NO:82);
- 25 g) AMB 2-4.1 (SEQ ID NO:90);
- h) AMB 2-3.1 (SEQ ID NO:91);
- i) AMB 2-5.1 (SEQ ID NO:92);
- j) AMB 2-6.1 (SEQ ID NO:93);
- k) AMB 2-2.1 (SEQ ID NO:94);
- 30 l) AMB 2-1.1 (SEQ ID NO:95);
- m) RAE 70.1-1 (SEQ ID NO:43);
- n) AMB 2-7.1 (SEQ ID NO:96);
- o) AMB 2-8.1 (SEQ ID NO:97);
- p) AMB 2-9.1 (SEQ ID NO:98);
- 35 q) AMB 2-10.1 (SEQ ID NO:99);
- r) AMB 2-11.1 (SEQ ID NO:100);
- s) RAE 70.1 (SEQ ID NO:44);
- t) AMB 3-4.1 (SEQ ID NO:103);
- u) AMB 3-5.1 (SEQ ID NO:102);

- 5                   v)     AMB 3-3.1 (SEQ ID NO:104);  
                  w)     AMB 3-2.1 (SEQ ID NO:105);  
                  x)     AMB 3-1.1 (SEQ ID NO:106);  
                  y)     AMB 4-8.1 (SEQ ID NO:109);  
                  z)     AMB 4-9.1 (SEQ ID NO:110);  
                  a')    AMB 4-6.1 (SEQ ID NO:111);  
                  b')    AMB 4-5.1 (SEQ ID NO:112);  
                  c')    AMB 4-3.1 (SEQ ID NO:113);  
                  d')    AMB 4-2.1 (SEQ ID NO:114); and  
10                  e')    AMB 4-1.1 (SEQ ID NO:115).

23.     An isolated nucleic acid having a sequence encoding all or a portion of a peptide of claim 20, or the functional equivalent of said nucleic acid sequence.

- 15           24.     An isolated peptide produced in a host cell transformed with the nucleic acid of claim 23.

25.     All or a portion of an isolated peptide of *Amb a* I.3, said peptide or portion thereof comprising at least one T cell epitope of *Amb a* I.3, said peptide comprising an amino  
20     acid sequence selected from the group consisting of:

- a)     AMB 1-4.15 (SEQ ID NO:88);  
                  b)     AMB 1-2.15 (SEQ ID NO:89);  
                  c)     AMB 2-1.15 (SEQ ID NO:101);  
25                  d)     AMB 3-4.15 (SEQ ID NO:107);  
                  e)     AMB 3-1.15 (SEQ ID NO:108); and  
                  f)     AMB 4-3.15 (SEQ ID NO:116).

30           26.     A therapeutic composition comprising all or a portion of an isolated peptide of claim 1 and a pharmaceutically acceptable carrier or diluent.

27.     A method of treating sensitivity to ragweed pollen in an individual, comprising administering to the individual a therapeutically effective amount of the composition of claim 26.

35           28.     A method of treating sensitivity to ragweed pollen in an individual, comprising administering sequentially to the individual a therapeutically effective amount of two different compositions of claim 26.

29. A therapeutic composition comprising all or a portion of an isolated peptide of claim 8 and a pharmaceutically acceptable carrier or diluent.

5 30. A therapeutic composition comprising all or a portion of an isolated peptide of claim 19 and a pharmaceutically acceptable carrier or diluent.

31. A therapeutic composition comprising all or a portion of an isolated peptide of claim 20 and a pharmaceutically acceptable carrier or diluent.

10 32. A method of treating sensitivity to ragweed pollen in an individual, comprising administering to the individual a therapeutically effective amount of the composition of claim 31.

15 33. A method of detecting sensitivity to ragweed pollen in an individual, comprising combining a blood sample obtained from the individual with all or a portion of peptide of claim 1, under conditions appropriate for binding of blood components with the peptide, and determining the extent to which such binding occurs as indicative of sensitivity in the individual to ragweed pollen.

20 34. A method of claim 33 wherein the extent to which binding occurs is determined by assessing T cell function, T cell proliferation or a combination thereof.

25 35. A method of detecting sensitivity to ragweed pollen in an individual, comprising combining a blood sample obtained from the individual with a peptide of claim 20, under conditions appropriate for binding of blood components with the peptide, and determining the extent to which such binding occurs as indicative of sensitivity in the individual to ragweed pollen.

30 36. A method of claim 35 wherein the extent to which binding occurs is determined by assessing T cell function, T cell proliferation or a combination thereof.

35 37. A composition comprising at least two peptides, said peptides each comprising at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia*, said peptides derived from the same or from different protein allergens of *Ambrosia artemisiifolia*, said composition selected from the group consisting of:

- a) AMB 4-6.1 and RAE 70.1 (SEQ ID NO:111 and SEQ ID NO:44);
- b) AMB 4-6.1 and AMB 2-5.1 (SEQ ID NO:111 and SEQ ID NO:92);
- c) AMB 4-9.1 and AMB 2-5.1 (SEQ ID NO:110 and SEQ ID NO:92);

- 5
- d) AMB 4-9.1 and RAE 70.1 (SEQ ID NO:110 and SEQ ID NO:44);
- e) AMB 4-6.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:92 and SEQ ID NO:86);
- f) AMB 4-9.1, AMB 2-5.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:92 and SEQ ID NO:86);
- 10
- g) AMB 4-6.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:111, SEQ ID NO:44 and SEQ ID NO:86);
- h) AMB 4-9.1, RAE 70.1 and AMB 1-2.1 (SEQ ID NO:110, SEQ ID NO:44 and SEQ ID NO:86);
- i) AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102);
- j) AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:102);
- 15
- k) AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102);
- l) AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-5.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:102);
- 20
- m) AMB 4-6.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103);
- n) AMB 4-9.1, RAE 70.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:44, SEQ ID NO:86 and SEQ ID NO:103);
- 25
- o) AMB 4-6.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:111, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103);
- p) AMB 4-9.1, AMB 2-5.1, AMB 1-2.1 and AMB 3-4.1 (SEQ ID NO:110, SEQ ID NO:92, SEQ ID NO:86 and SEQ ID NO:103);
- 30
- q') AMB 2-1.15 and AMB 4-3.15 (SEQ ID NO:101, and SEQ ID NO:116);
- r') AMB 1-2.15, AMB 2-1.15 and  
AMB 4-3.15 (SEQ ID NO:89, SEQ ID NO:101, and SEQ ID NO:116);  
and
- 35
- s') AMB 1-2.15, AMB 2-1.15,  
AMB 4-3.15 and AMB 3-4.15 (SEQ ID NO:89, SEQ ID NO:101, SEQ ID NO:116 and SEQ ID NO:107).

38. A therapeutic composition comprising a pharmaceutically acceptable carrier or diluent and at least two peptides, said peptides each comprising at least one T cell epitope of a protein allergen of *Ambrosia artemisiifolia*, said peptides derived from the same or from different protein allergens of *Ambrosia artemisiifolia*.

5

39. A method of treating sensitivity to ragweed pollen in an individual, comprising administering to the individual a therapeutically effective amount of the composition of claim 38.

10

40. A method of treating sensitivity to ragweed pollen in an individual, comprising administering to the individual a therapeutically effective amount of a therapeutic composition comprising a pharmaceutically acceptable carrier or diluent and a protein allergen selected from the group consisting of recombinant *Amb a* I.1, recombinant *Amb a* I.2, recombinant *Amb a* I.3, recombinant *Amb a* I.4 and recombinant *Amb a* II, said protein  
15 allergen which binds immunoglobulin E to a substantially lesser extent than the corresponding native protein allergen.

20

41. An isolated peptide comprising at least two different regions, each region comprising at least one T cell epitope of *Amb a* I.1, said regions each comprising all or a portion of an amino acid sequence selected from the group consisting of: amino acid residues 48-107 (SEQ ID NO:117); amino acid residues 171-216 (SEQ ID NO:118); amino acid residues 278-322 (SEQ ID NO:119); and amino acid residues 331-377 (SEQ ID NO:120).

25

42. An isolated peptide comprising at least two different regions, each region comprising at least one T cell epitope of *Amb a* I.1, said regions each comprising all or a portion of an amino acid sequence selected from the group consisting of: amino acid residues 57-101 (SEQ ID NO:121); amino acid residues 182-216 (SEQ ID NO:122); amino acid residues 280-322 (SEQ ID NO:123); and amino acid residues 342-377 (SEQ ID  
30 NO:124).

35

43. An isolated peptide of claim 41 wherein the regions comprise amino acid residues 171-216 (SEQ ID NO:118) of *Amb a* I.1 and amino acid residues 331-377 (SEQ ID NO:120) of *Amb a* I.1.

44. An isolated peptide of *Amb a* I.1 comprising at least one T cell epitope of *Amb a* I.1, said peptide comprising all or a portion of an amino acid sequence selected from the group consisting of: amino acid residues 48-107 (SEQ ID NO:117); amino acid residues



171-216 (SEQ ID NO:118); amino acid residues 278-322 (SEQ ID NO:119); and amino acid residues 331-377 (SEQ ID NO:120).

5 45. A therapeutic composition comprising at least one peptide of *Amb a* I.1 and a pharmaceutically acceptable carrier or diluent, said at least one peptide comprising a sufficient percentage of the T cell epitopes of *Amb a* I.1 such that upon administration of the peptide to an individual sensitive to ragweed pollen, T cells of the individual are tolerized to *Amb a* I.1.

10 46. A method of treating sensitivity to ragweed pollen in an individual, comprising a therapeutically effective amount of a composition of claim 45.

15 47. A method of preventing sensitivity to ragweed pollen in an individual, comprising administering to the individual a therapeutically effective amount of a composition of claim 45.

48. A composition comprising at least two peptides, said peptides each comprising at least one T cell epitope of *Amb a* I.1, said composition selected from the group consisting of:

- 20 a) AMB 1-2.1 and AMB 4-9.1 (SEQ ID NO:86 and SEQ ID NO:110);  
b) AMB 1-2.1, AMB 2-38.1 and AMB 4-9.1 (SEQ ID NO:86, SEQ ID NO:141, and SEQ ID NO:110);  
c) AMB 1-2.1, AMB 2-38.1, AMB 4-9.1 and AMB 2-4.1 (SEQ ID NO:86, SEQ ID NO:141, SEQ ID NO:110, and SEQ ID NO:90);  
25 d) AMB 1-2.1, AMB 2-38.1, AMB 4-9.1, AMB 2-4.1, and AMB 3-5.1 (SEQ ID NO:86, SEQ ID NO:141, SEQ ID NO:110, SEQ ID NO:90, and SEQ ID NO:102);  
e) AMB 1-2.1, AMB 2-36.1 and AMB 4-9.1 (SEQ ID NO:86, SEQ ID NO:139, and SEQ ID NO:110);  
30 f) AMB 1-2.1, AMB 2-36.1, AMB 4-9.1 and AMB 2-4.1 (SEQ ID NO:86, SEQ ID NO:139, SEQ ID NO:110, and SEQ ID NO:90); and  
g) AMB 1-2.1, AMB 2-36.1, AMB 4-9.1, AMB 2-4.1, and AMB 3-5.1 (SEQ ID NO:86, SEQ ID NO:139, SEQ ID NO:110, SEQ ID NO:90, and SEQ ID NO:102).  
35

49. A peptide of claim 8 in which at least one amino acid residue not essential for T cell receptor interaction has been substituted with another amino acid residue.

50. A peptide of claim 49 wherein said at least one amino acid residue not essential to T cell receptor interaction has been substituted with an amino acid residue selected from the group consisting of a methyl amino acid, alanine and glutamic acid.

5

51. A peptide of claim 8 wherein at least one amino acid residue essential for T cell receptor interaction has been substituted with a similar amino acid residue.

10

52. A peptide of claim 8 which has been modified by a process comprising the steps of:

15

- a) determining the amino acid residues of said peptide essential to T cell receptor recognition;
- b) substituting at least one amino acid residue not essential to T cell receptor recognition with an amino acid residue, selected from the group consisting of a methyl amino acid, alanine and glutamic acid; and optionally
- c) substituting at least one amino acid residue essential for T cell interaction with a similar amino acid residue.

20

53. A peptide of claim 8 modified by a process comprising the steps of:

25

- a) determining the amino acid residues of said peptide which are essential to T cell receptor recognition;
- b) substituting at least one amino acid residue essential for T cell interaction with a similar amino acid residue; and optionally
- c) substituting at least one amino acid residue not essential to T cell receptor interaction with an amino acid residue selected from the group consisting of methyl amino acid, alanine and glutamic acid.

30

54. A method for determining in an individual the presence of immunoglobulin E specific for a ragweed protein allergen, and the ability of T cells of the individual to respond to T cell epitope(s) of ragweed protein allergen, comprising the steps of:

35

- a) administering to an individual an Immediate Type Hypersensitivity test utilizing *Amb a* I.1 or a portion thereof which binds immunoglobulin E specific for the human T cell reactive feline protein;

- b) determining whether a specific Immediate Type Hypersensitivity reaction occurs;

- c) administering to an individual prior to, simultaneously with, or subsequent to administration of the Immediate Type Hypersensitivity test in step (a), a Delayed Type

Hypersensitivity test utilizing a peptide derived from *Amb a* 1.1 which has human T cell stimulating activity and which does not bind immunoglobulin E specific for *Amb a* 1.1, or if binding to said immunoglobulin E occurs, such binding does not result in release of mediators from mast cells or basophils in a substantial percentage of a population of individuals sensitive to the human T cell reactive feline protein; and

d) determining whether a specific Delayed Type Hypersensitivity reaction occurs.

55. An antibody specifically reactive with a peptide of claim 1.
56. The antibody of claim 55 which is a monoclonal antibody.
57. A T cell clone specifically reactive with a peptide of claim 1.
58. A soluble T cell receptor specifically reactive with a peptide of claim 1.
59. An antibody specifically reactive with the T cell receptor of claim 58.
60. A multi-peptide formulation suitable for pharmaceutical administration, comprising: at least two peptides derived from a ragweed pollen protein allergen or a combination thereof, having human T cell stimulating activity in an *in vitro* T cell proliferation assay and soluble at a physiologically acceptable predetermined pH and an excipient.
61. The multi-peptide formulation of claim 60 said at least two peptides are lyophilized.
62. The multi-peptide formulation of claim 60, wherein the excipient comprises sodium phosphate.
63. The multi-peptide formulation of claim 60, wherein the excipient comprises mannitol.
64. A multi-peptide formulation comprising peptides Amb 1-2.1 (SEQ ID NO:86) Amb 2-36.1 (SEQ ID NO: 139), and Amb 4-9.1 (SEQ ID NO: 110).
65. The multi-peptide formulation of claim 64 further comprising sodium phosphate and mannitol.

66. The multipeptide formulation of claim 64 wherein said formulation is in lyophilized form.

5 67. A therapeutic composition comprising at least one peptide of *Amb a* I.1, , at least one said peptide containing a sufficient percentage of the T cell epitopes present in ragweed pollen protein allergen to induce T cell nonresponsiveness in a substantial percentage of a population of individuals sensitive to ragweed pollen protein allergen.

10 — 68. A therapeutic composition comprising at least one peptide of claim 1, at least one said peptide comprising a sufficient percent of the T cell epitopes present in *Amb a* I.1 to induce T cell nonresponsiveness in a substantial percent of the population of individuals sensitive to ragweed protein allergen.

15 69. A method for treating sensitivity to a ragweed pollen protein allergen in a subject sensitive to said allergen comprising orally administering at least one isolated and purified ragweed pollen protein allergen or any portion thereof.

20 70. The composition of claim 48 comprising AMB 1-2.1, AMB 2-36.1 and AMB 4-9.1 (SEQ ID NO:86, SEQ ID NO:139, and SEQ ID NO:110)

71. A method of treating sensitivity to ragweed pollen protein allergen in an individual comprising administering subcutaneously, in non-immunogenic form, a therapeutically effective amount of the multipeptide formulation of claim 65.

1 / 39




FIG. 1

2/39




Fig. 2

3/39




Fig. 3





Fig. 4A








Fig. 5A




Fig. 5B

6/39




Fig. 5C




Fig. 5D

7/39




Fig. 5E




Fig. 5F

8/39




Fig. 6A




Fig. 6B

SUBSTITUTE SHEET (RULE 61)

9/39




Fig. 6C




Fig. 6D

10/39




Fig. 6E




Fig. 6F

11/39

RAE 67.1(8-34)  
RAE 67.15(8-34)  
RAE 57.1(22-51)35,42del  
RAE 24.E(38-61)42S44K  
RAE 24.1(38-61)42del.  
RAE 22.E(48-75)  
RAE 22.E-1(57-75)  
RAE 3.D(58-67C)  
RAE 3.1(58-67)  
RAE 22.E-2(63-75)  
RAE 5.D(67-83)  
RAE 6.D(75-92)90S  
RAE 6.1(75-92)  
RAE 7.D(84-100)90S  
RAE 7.D-1(84-100)  
RAE 40.1-6(88-98)  
RAE 40.1-5(88-101)  
RAE 40.1-4(88-104)  
RAE 40.D(88-107)90S  
RAE 40.1(88-107)  
RAE 61.1(101-129)  
RAE 80.1(118-137)  
RAE 45.1(129-147)  
RAE 45.15(129-147)

YILYFTLALVTLLQPVRSAEDLQEILP  
YILYFTLALVALLOPVRSAEGVGEILP  
PVRSAEDLQEILP\*VNETRR\*LTSGAYNI  
ETRRSLKTSAYNIIDGCWRGKAD  
ETRR\*LTSGAYNIIDGCWRGKAD  
AYNIIDGCWRGKADWAENRRKALADCAQG  
RGKADWAENRRKALADCAQG  
GKADWAENRC  
GKADWAENRK  
AENRRKALADCAQG  
KALADCAQGGFGKGTGG  
GFGKGTGGKDGDIYSVT  
GFGKGTGGKDGDIYTVT  
KDGDIYSVTSELDDDDVA  
KDGDIYTVTSELDDDDVA  
IYTVTSELDDDD  
IYTVTSELDDDDVAN  
IYTVTSELDDDDVANPKE  
IYSVTSELDDDDVANPKEGTL  
IYTVTSELDDDDVANPKEGTL  
NPKEGTLRFGAQNRPLWIIFERDMVIRL  
WIIFERDMVIRLDKEMVVNS  
LDKEMVVNSDKTIDGRGAK  
LNQELVVNSDKTIDGRGVK

Fig. 7

12/39

RAE 45.15(129-147)	LNQELVVNSDKTIDGRGVK
RAE 75.1(138-163)	DKTIDGRGAKVEIINAGFTLNGVKNV
RAE 62.1(154-180)	GFTLNGVKNV I IHNNMHVDVKNPGL
RAE 69.1(171-198)	HDVKVNPGLIKSNDGPAAPRAGSDGDA
RAE 69.1-1(176-198)	NPGGLIKSNDGPAAPRAGSDGDA
RAE 69.1-2(181-198)	IKSNDGPAAPRAGSDGDA
RAE 69.1-3(186-198)	GPAAPRAGSDGDA
RAE 70.1-3(189-201)	APRAGSDGDAISI
RAE 70.1-2(189-206)	APRAGSDGDAISISGSSQ
RAE 70.1-1(189-211)	APRAGSDGDAISISGSSQIWDH
RAE 70.1(189-216)	APRAGSDGDAISISGSSQIWDHCSLSK
RAE 71.1(208-235)	WIDHCSLSKSVGLVDAKLGTTRLTVSN
RAE 65.1(226-252)	LGTRLTVSNSLFTQHGFVLLFGAGDE
RAE 63.1(243-270)	FVLLFGAGDENIEDRCMLATVAFNTFTD
RAE 76.1(260-277)	LATVAFNTFTDNVDQRMP
RAE 76.6(260-277)	LATVAFNMFTDHVDQRMP
RAE 27.1(268-287)	FTDNVDQRMPRCRHGFFQVV
RAE 27.15(268-287)	FTDNVDQRMPRCRFGFFQVV
RAE 66.1(278-305)	RCRHGFFQVVNNNYDKWGSYAIG
RAE 66.1-1(281-305)	HGFFQVVNNNYDKWGSYAIGGSASP
RAE 66.1-2(284-305)	FQVVNNNYDKWGSYAIGGSASP
RAE 66.1-3(287-305)	VNNNYDKWGSYAIGGSASP
RAE 64.1-3(295-306)	GSYAIGGSASPT

Fig. 7 cont



13/39

RAE 64.1-2(295-310) GSYAIGGSASPTILSQ  
RAE 64.1-1(295-314) GSYAIGGSASPTILSQGNRF  
RAE 64.1(295-322) GSYAIGGSASPTILSQGNRFCAPDERSK  
RAE 73.1(313-340) RFCAPDERSKKNVLGRHGEAAESMKWN  
RAE 74.1(331-356) EAAAESMKWNWRTNKDVLENGAIFVA  
RAE 74.1-1(335-357) ESMKWNWRTNKDVLENGAIFVAS  
RAE 29.1(338E-357)339F340G EFGWRTNKDVLENGAIFVAS  
RAE 29.1-1(338-357) KWNWRTNKDVLENGAIFVAS  
RAE 28+29(338-367) KWNWRTNKDVLENGAIFVASGVDPVLTPEQ  
RAE 29.1-2(341-357) WRTNKDVLENGAIFVAS  
RAE 29.1-3(344-357) NKDVLENGAIFVAS  
RAE 29.1-4(347-357) VLENGAIFVAS  
RAE 28.1-3(348-358) LENGAIFVASG  
RAE 28.1-2(348-361) LENGAIFVASGVDP  
RAE 28.1-1(348-364) LENGAIFVASGVDPVLT  
RAE 28.1(348-367) LENGAIFVASGVDPVLTPEQ  
RAE 20.1(358-377) GVDPVLTPEQSAGMIPAEPG  
RAE 20.1-3(358-366) GVDPVLTPE  
RAE 20.1-2(358-369) GVDPVLTPEQSA  
RAE 20.1-1(358-373) GVDPVLTPEQSAGMIP  
RAE 21.1(368-387) SAGMIPAEPGESALSLTSSA  
RAE 17.1(375-392) EPGESALSLTSSAGVLSC  
RAE 55.1(384-398) TSSAGVLSCQPGAPC

Fig. 7 cont

14/39




Fig. 8

15/39

RAE 22.E(48-75)  
RAE 22.E-1(57-75)  
RAE 22.E-2(63-75)  
RAE 5.D(67-83)  
RAE 6.D(75-92)90S  
RAE 7.D(84-100)90S  
RAE 40.1(88-107)  
RAE 40.1-4(88-104)  
RAE 40.1-5(88-101)  
RAE 40.1-6(88-98)  
RAE 69.1(171-198)  
RAE 69.1-1(176-198)  
RAE 69.1-2(181-198)  
RAE 69.1-3(186-198)  
RAE 70.1(189-216)  
RAE 70.1-1(189-211)  
RAE 70.1-2(189-206)  
RAE 70.1-3(189-201)  
RAE 66.1(278-305)  
RAE 66.1-1(281-305)  
RAE 66.1-2(284-305)

AYNIIDGWRGKADWAENRKALADCAQG  
RGKADWAENRKALADCAQG  
AENRKALADCAQG  
KALADCAQGFKGKTVGG  
GFGKGTGGKDDIYSVT  
KGDIIYSVTSELDDDDVA  
IYTVTSELDDDDVANPKEGTL  
IYTVTSELDDDDVANPKE  
IYTVTSELDDDDVAN  
IYTVTSELDDDD  
HDVKVNPGLIKSNDGPAAPRAGSDGDA  
NPGGLIKSNDGPAAPRAGSDGDA  
IKSNDGPAAPRAGSDGDA  
GPAAPRAGSDGDA  
APRAGSDGDAISISGSSQIWDHCSLSK  
APRAGSDGDAISISGSSQIWDH  
APRAGSDGDAISISGSSQ  
APRAGSDGDAISI  
RCRHGFFQVNNNNYDKWGSYAIGGSASP  
HGFFQVNNNNYDKWGSYAIGGSASP  
FQVNNNNYDKWGSYAIGGSASP

Fig. 9


16/39

RAE 66.1-3(287-305)  
RAE 64.1(295-322)  
RAE 64.1-1(295-314)  
RAE 64.1-2(295-310)  
RAE 64.1-3(295-306)  
RAE 74.1(331-356)  
RAE 74.1-1(335-357)  
RAE 29.1-1(338-357)  
RAE 28+29(338-367)  
RAE 29.1-2(341-357)  
RAE 29.1-3(344-357)  
RAE 29.1-4(347-357)  
RAE 28.1-3(348-358)  
RAE 28.1-2(348-361)  
RAE 28.1-1(348-364)  
RAE 28.1(348-367)  
RAE 20.1(358-377)  
RAE 20.1-1(358-373)  
RAE 20.1-2(358-369)  
RAE 20.1-3(358-366)

VNNYDKWGSYAIGGSASP  
GSYAIGGSASPTILSQNRFCAPDERSK  
GSYAIGGSASPTILSQGNRF  
GSYAIGGSASPTILSQ  
GSYAIGGSASPT  
EAAESMKWNWRTNKDVLENGAIFVA  
ESMKWNWRTNKDVLENGAIFVAS  
KWNWRTNKDVLENGAIFVAS  
KWNWRTNKDVLENGAIFVASGVDPVLTPEQ  
WRTNKDVLENGAIFVAS  
NKDVLENGAIFVAS  
VLENGAIFVAS  
LENGAIFVASG  
LENGAIFVASGVDP  
LENGAIFVASGVDPVLT  
LENGAIFVASGVDPVLTPEQ  
GVDPVLTPEQSAGMIPAEFG  
GVDPVLTPEQSAGMIP  
GVDPVLTPEQSA  
GVDPVLTPE

Fig. 9 cont


17/39



AMB A I.1 PEPTIDES

Fig. 10


18/39



AMB A I.I PEPTIDES

Fig. 11


19/39



AMB A.I.1 PEPTIDES

Fig. 12

20/39



AMB A1.1 PEPTIDES

Fig. 13



21/39

AMB 1-6.1(57-84)	RGKADWAENRKALADCAQGFVGKGTGGK
AMB 1-5.1(63-84)	AENRKALADCAQGFVGKGTGGK
AMB 1-4.1(63-87)	AENRKALADCAQGFVGKGTGGKDG
AMB 1-1.1(75-101)	GFGKGTGGKGDIIYTVTSELDDDDVAN
AMB 1-2.1(78-101)	KGTGGKGDIIYTVTSELDDDDVAN
AMB 1-3.1(83-101)	GKGDIIYTVTSELDDDDVAN
AMB 1-4.15(63-87)	ENNRQALADCAQGFVGKGTGGKWD
AMB 1-2.15(78-101)	KGTGGKWDVYTVTSNLDDDDVAN
AMB 2-4.1(182-203)	KSNDGPAAPRAGSDGDAISISG
AMB 2-3.1(182-206)	KSNDGPAAPRAGSDGDAISISGSSQ
AMB 2-5.1(182-211)	KSNDGPAAPRAGSDGDAISISGSSQIWDH
AMB 2-6.1(182-216)	KSNDGPAAPRAGSDGDAISISGSSQIWDHCSLSK
AMB 2-2.1(186-203)	GPAAPRAGSDGDAISISG
AMB 2-1.1(186-206)	GPAAPRAGSDGDAISISGSSQ
AE 70.1-1(189-211)	APRAGSDGDAISISGSSQIWDH
MB 2-7.1KK(189-211)KK	KKAPRAGSDGDAISISGSSQIWDHKK
AE 70.1(189-216)	APRAGSDGDAISISGSSQIWDHCSLSK
MB 2-8.1KK(189-216)K	KKAPRAGSDGDAISISGSSQIWDHCSLSK
MB 2-9.1KK(189-216)K, S212	KKAPRAGSDGDAISISGSSQIWDHSSLSK
MB 2-10.1KK(189-216)K, L212	KKAPRAGSDGDAISISGSSQIWDHLSLSK
MB2-11.1KK(189-216)K, E212	KKAPRAGSDGDAISISGSSQIWDHESLSK

Fig. 14

22/39

AMB 2-1.15(186-206)	GPILRQASDGTINVAGSSQ
AMB 3-5.1(280-306)	RHGEFFQVNNNYDKWGSYAIGGSASPT
AMB 3-4.1(283-306)	FFQVNNNYDKWGSYAIGGSASPT
AMB 3-3.1(295-318)	GSYAIGGSASPTILSQGNRFCAPD
AMB 3-2.1(295-320)	GSYAIGGSASPTILSQGNRFCAPDER
AMB 3-1.1(295-322)	GSYAIGGSASPTILSQGNRFCAPDERSK
AMB 3-4.15(283-306)	FFQVNNNYDRWGTYAIGGSAPT
AMB 3-1.15(295-322)	GTYAIGGSAPTILCQGNRFLAPDDQIK
AMB 4-8.1(345-373)	KDVLENGAIFVASGVDPVLTPEQSAGMIP
AMB 4-9.1(345-370)	KDVLENGAIFVASGVDPVLTPEQSAG
AMB 4-6.1(347-377)	VLENGAIFVASGVDPVLTPEQSAGMIPAEPG
AMB 4-5.1(347-375)	VLENGAIFVASGVDPVLTPEQSAGMIPAE
AMB 4-3.1(347-373)	VLENGAIFVASGVDPVLTPEQSAGMIP
AMB 4-2.1(347-370)	VLENGAIFVASGVDPVLTPEQSAG
AMB 4-1.1(347-367)	VLENGAIFVASGVDPVLTPEQ
AMB 4-3.15(347-373)	LENGAIFVTSGSDPVLTVPVQSAGMIP

Fig. 14 cont

23/39




Fig. 15

24/39




Fig. 16

25/39




Fig. 17

26/39




Fig. 18




Fig. 19

28/39




Fig. 20



29/39




Fig. 21




Fig. 22

31/39




Fig. 23

32/39




Fig. 24

33/39

NAME	SEQUENCE	CRUDE YIELD
ORIGINAL	-SDGDAISISGSSQIWIDHCSLSKS	-----
Amb2-18.1	SGSSQIWIDHSSLKS	79%
Amb2-19.1	SISGSSQIWIDHSSLKS	67%
Amb2-20.1	DAISISGSSQIWIDHSSLKS	80%
Amb2-21.1	SDGDAISISGSSQIWIDHSSLKS	87%
Amb2-22.1	SESGSSQIWIDHSSLKS	84%
Amb2-23.1	SKSGSSQIWIDHSSLKS	87%
Amb2-26.1	DKSGSSQIWIDHSSLSKE	
Amb2-28.1	DKSISGSSQIWIDHSSLSKE	98%
Amb2-30.1	DKESGSSQIWIDHSSLSKE	
Amb2-32.1	DKSGSSQIWIDHSSLSKK	85%
Amb2-33.1	DKESGSSQIWIDHSSLSKEK	107%
Amb2-34.1	SGSSQIAIDHSSLKS	
Amb2-35.1	DKESGSSQIAIDHSSLSKSE	
Amb2-36.1	DKESGSSQIWIDHSSLSKEKD	
Amb2-37.1	SQIWIDHSSLKS	
Amb2-38.1	DKESISGSSQIWIDHSSLSKEKD	

Fig. 25

34/39




Fig. 26




Fig. 27

36/39




Fig. 28



37/39




Fig. 29

38/39




Fig. 30

39/39




FIGURE 31

# INTERNATIONAL SEARCH REPORT

Inter Application No  
PCT/US 95/14362

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/29 C07K14/415 A61K39/36 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 21321 (IMMULOGIC PHARMACEUTICAL CORPORATION) 28 October 1993 see the whole document -----	1-71

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\* document member of the same patent family

Date of the actual completion of the international search

14 March 1996

Date of mailing of the international search report

26. 03.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Cupido, M

# INTERNATIONAL SEARCH REPORT

ional application No.  
PCT/US95/ 14362

## B x I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 27,28,32-36,39,40,46,47,54,69 and 71 are directed to a method of treatment or diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compositions.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Inte Application No  
PCT/US 95/14362

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9321321	28-10-93	AU-B- 4046293	18-11-93
		CA-A- 2117779	28-10-93
		EP-A- 0636180	01-02-95
		FI-A- 944697	07-12-94
		JP-T- 7507924	07-09-95
		NO-A- 943811	06-12-94
		ZA-A- 9302530	28-10-93
-----			